# **PCT**

# WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



# INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7: (11) International Publication Number: WO 00/28061
C12N 15/86, 15/35, 5/10, A61K 48/00 A2 (43) International Publication Date: 18 May 2000 (18.05.00)

US

(21) International Application Number:

PCT/US99/25694

(22) International Filing Date:

2 November 1999 (02.11.99)

(30) Priority Data: 60/107,114

5 November 1998 (05.11.98)

(71) Applicant (for all designated States except US): THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Suite 300, 3700 Market Street, Philadelphia, PA 19104-3147 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): WILSON, James, M. [US/US]; 1350 N. Avignon Drive, Gladwyne, PA 19035 (US). XIAO, Weidong [CN/US]; Apartment P4, 155 Washington Lane, Jenkintown, PA 19046 (US).

(74) Agents: KODROFF, Cathy, A. et al.; Howson & Howson, Spring House Corporate Center, P.O. Box 457, Spring House, PA 19477 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), Buropean patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

#### Published

Without international search report and to be republished upon receipt of that report.

(54) Title: ADENO-ASSOCIATED VIRUS SEROTYPE 1 NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

### (57) Abstract

The nucleic acid sequences of adeno-associated virus (AAV) serotype 1 are provided, as are vectors and host cells containing these sequences and functional fragments thereof. Also provided are methods of delivering genes via AAV-1 derived vectors.

2

# FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	E8	Spain	LS	Lesotho	81	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	8K	Slovakia
AT	Austria	FR	Presce	LU	Luxembourg	SN	Senegal
ΑU	Australia	GA	Gabon	LV	Latvia	<b>8</b> Z	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GB	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Chana	MG	Madagascar	TJ	Tajikistan -
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobego
BJ	Benin	IB	freland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	(S	Iceland	MW	Malewi	US	United States of Americ
CA	Canada	ſŦ	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NB	Niger	VN	Vict Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan -	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	ΚZ	Kazakatan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	Ц	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
BB.	Estonia	LR	Liberia	SG	Singapore		

WO 00/28061 PCT/US99/25694

# ADENO-ASSOCIATED VIRUS SEROTYPE I NUCLEIC ACID SEQUENCES, VECTORS AND HOST CELLS CONTAINING SAME

This work was supported by the National Institutes of Health, grant no. P30 DK47757-06 and PO1 HD32649-04. The US government may have certain rights in this invention.

#### Field of the Invention

5

This invention relates generally to viral vector, and more particularly, to recombinant viral vectors useful for gene delivery.

### **Background of the Invention**

Adeno-associated viruses are small, single-stranded DNA viruses which 10 require helper virus to facilitate efficient replication [K.I. Berns, Parvoviridae: the viruses and their replication, p. 1007-1041, in F.N. Fields et al., Fundamental virology, 3rd ed., vol. 2, (Lippencott-Raven Publishers, Philadelphia, PA) (1995)]. The 4.7 kb genome of AAV is characterized by two inverted terminal repeats (ITR) and two open reading frames which encode the Rep proteins and Cap proteins, 15 respectively. The Rep reading frame encodes four proteins of molecular weight 78 kD, 68 kD, 52 kD and 40 kD. These proteins function mainly in regulating AAV replication and integration of the AAV into a host cell's chromosomes. The Cap reading frame encodes three structural proteins in molecular weight 85 kD (VP 1), 72 20 kD (VP2) and 61 kD (VP3) [Berns, cited above]. More than 80% of total proteins in AAV virion comprise VP3. The two ITRs are the only cis elements essential for AAV replication, packaging and integration. There are two conformations of AAV ITRs called "flip" and "flop". These differences in conformation originated from the replication model of adeno-associated virus which use the ITR to initiate and reinitiate 25 the replication [R.O. Snyder et al., J. Virol., 67:6096-6104 (1993); K.I. Berns, Microbiological Reviews, 54:316-329 (1990)].

AAVs have been found in many animal species, including primates, canine, fowl and human [F.A. Murphy et al., "The Classification and Nomenclature of Viruses: Sixth Report of the International Committee on Taxonomy of Viruses",

WO 00/28061 PCT/US99/25694

2

Archives of Virology, (Springer-Verlag, Vienna) (1995)]. In addition to five known primate AAVs (AAV-1 to AAV-5), AAV-6, another serotype closely related to AAV-2 and AAV-1 has also been isolated [E. A. Rutledge et al., J. Virol., 72:309-319 (1998)]. Among all known AAV serotypes, AAV-2 is perhaps the most well-characterized serotype, because its infectious clone was the first made [R.J. Samulski et al., Proc. Natl. Acad. Sci. USA, 79:2077-2081 (1982)]. Subsequently, the full sequences for AAV-3A, AAV-3B, AAV-4 and AAV-6 have also been determined [Rutledge, cited above; J.A.Chiorini et al., J. Virol., 71:6823-6833 (1997); S. Muramatsu et al., Virol., 221:208-217 (1996)]. Generally, all AAVs share more than 80% homology in nucleotide sequence.

5

10

15

20

25

A number of unique properties make AAV a promising vector for human gene therapy [Muzyczka, Current Topics in Microbiology and Immunology, 158:97-129 (1992)]. Unlike other viral vectors, AAVs have not been shown to be associated with any known human disease and are generally not considered pathogenic. Wild type AAV is capable of integrating into host chromosomes in a site specific manner [R. M. Kotin et al., Proc, Natl. Acad, Sci. USA, 87:2211-2215 (1990)- R.J. Samulski, EMBO J., 10(12):3941-3950 (1991)]. Recombinant AAV vectors can integrate into tissue cultured cells in chromosome 19 if the rep proteins are supplied in trans [C. Balague et al., J. Virol., 71:3299-3306 (1997); R. T. Surosky et al., J. Virol., 71:7951-7959 (1997)]. The integrated genomes of AAV have been shown to allow long term gene expression in a number of tissues, including, muscle, liver, and brain [K. J. Fisher, Nature Med., 3(3):306-312 (1997); R. 0. Snyder et al., Nature Genetics, 16:270-276 (1997); X. Xiao et al., Experimental Neurology, 144:113-124 (1997); Xiao, J. Virol., 70(11):8098-8108 (1996)].

AAV-2 has been shown to be present in about 80-90% of the human population. Earlier studies showed that neutralizing antibodies for AAV-2 are prevalent [W. P. Parks et al., <u>J. Virol.</u>, <u>2</u>:716-722 (1970)]. The presence of such antibodies may significantly decrease the usefulness of AAV vectors based on AAV-2 despite its other merits. What are needed in the art are vectors characterized by the

WO 00/28061 PCT/US99/25694

3

advantages of AAV-2, including those described above, without the disadvantages, including the presence of neutralizing antibodies.

### Summary of the Invention

5

10

15

20

In one aspect, the invention provides an isolated AAV-1 nucleic acid molecule which is selected from among SEQ ID NO: 1, the strand complementary to SEQ ID NO: 1, and cDNA and RNA sequences complementary to SEQ ID NO: 1 and its complementary strand.

In another aspect, the present invention provides AAV ITR sequences, which include the 5' ITR sequences, nt 1 to 143 of SEQ ID NO: 1; the 3' ITR sequences, nt 4576 to 4718 of SEQ ID NO: 1, and fragments thereof.

In yet another aspect, the present invention provides a recombinant vector comprising an AAV-1 ITR and a selected transgene. Preferably, the vector comprises both the 5' and 3' AAV-1 ITRs between which the selected transgene is located.

In still another aspect, the invention provides a recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.

In a further aspect, the present invention provides a nucleic acid molecule encoding an AAV-1 rep coding region and an AAV-1 cap coding region.

In still another aspect, the present invention provides a host cell transduced with a recombinant viral vector of the invention. The invention further provides a host cell stably transduced with an AAV-1 P5 promoter of the invention.

In still a further aspect, the present invention provides a pharmaceutical composition comprising a carrier and a vector of the invention.

In yet another aspect, the present invention provides a method for AAV—
mediated delivery of a transgene to a host involving the step of delivering to a selected host a recombinant viral vector comprising a selected transgene under the control of sequences which direct expression thereof and an adeno-associated virus 1 (AAV-1) virion.

In another aspect, the invention provides a method for in vitro production of a selected gene product using a vector of the invention.

Other aspects and advantages of the invention will be readily apparent to one of skill in the art from the detailed description of the invention.

### 5 Brief Description of the Drawings

10

15

20

25

Figs. 1A-1C illustrate the alignment of nucleotides of AAV-1 [SEQ ID NO: 1], AAV-2 [SEQ ID NO: 18] and AAV-6 [SEQ ID NO: 19]. The alignment was done with MacVector 6.0. The full sequences of AAV-1 are shown in the top line. Nucleotides in AAV-2 and AAV-6 identical to AAV-1 are symbolized by "." and gaps by "-". Some of the conserved features among AAVs are marked in this figure. Note the 3' ITRs of AAV-1 and AAV-6 are shown in different orientations.

Fig. 2 illustrates the predicted secondary structure of AAV-1 ITR. The nucleotides in AAV-2 and AAV-6 are shown in italic and bold respectively.

Fig. 3A illustrates a hypothesis of how AAV-6 arose from the homologous recombination between AAV-1 and AAV-2. The major elements of AAV-1 are indicated in the graph. A region that is shared between AAV-1, AAV-2 and AAV-6 is shown in box with waved lines.

Fig. 3B is a detailed illustration of a 71 bp homologous region among AAV-1, AAV-2 and AAV-6. Nucleotides that differ among these serotypes are indicated by arrows.

Fig. 4A is a bar chart illustrating expression levels of human alpha 1 antitrypsin (α1AT) in serum following delivery of hAAT via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 4B is a bar chart illustrating expression levels of erythropoietin (epo) in serum following delivery of the epo gene via recombinant AAV-1 and recombinant AAV-2 viruses.

Fig. 5A is a bar chart illustrating expression levels of a1AT in liver following delivery of a1AT as described in Example 7.

10

20

25

Fig. 5B is a bar chart demonstrating expression levels of epo in liver following delivery of epo as described in Example 7.

Fig. 5C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of a1AT or epo to liver as described in Example 7.

Fig. 5D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of α1 AT or epo to liver as described in Example 7.

Fig. 6A is a bar chart illustrating expression levels of  $\alpha 1$  AT in muscle following delivery of  $\alpha 1$  AT as described in Example 7.

Fig. 6B is a bar chart demonstrating expression levels of epo in muscle following delivery of epo as described in Example 7.

Fig. 6C is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-1 following delivery of α1AT or epo to muscle as described in Example 7.

Fig. 6D is a bar chart demonstrating neutralizing antibodies (NAB) directed to AAV-2 following delivery of al AT or epo to muscle as described in Example 7.

#### 15 Detailed Description of the Invention

The present invention provides novel nucleic acid sequences for an adeno-associated virus of serotype 1 (AAV-1). Also provided are fragments of these AAV-1
sequences. Among particularly desirable AAV-1 fragments are the inverted terminal
repeat sequences (ITRs), rep and cap. Each of these fragments may be readily
utilized, e.g., as a cassette, in a variety of vector systems and host cells. Such
fragments may be used alone, in combination with other AAV-1 sequences or
fragments, or in combination with elements from other AAV or non-AAV viral
sequences. In one particularly desirable embodiment, a cassette may contain the
AAV-1 ITRs of the invention flanking a selected transgene. In another desirable
embodiment, a cassette may contain the AAV-1 rep and/or cap proteins, e.g., for use
in producing recombinant (rAAV) virus.

Thus, the AAV-1 sequences and fragments thereof are useful in production of rAAV, and are also useful as antisense delivery vectors, gene therapy vectors, or vaccine vectors. The invention further provides nucleic acid molecules, gene delivery

10

15

20

25

vectors, and host cells which contain the AAV-1 sequences of the invention. Also provided a novel methods of gene delivery using AAV vectors.

As described herein, the vectors of the invention containing the AAV-1 capsid proteins of the invention are particularly well suited for use in applications in which the neutralizing antibodies diminish the effectiveness of other AAV serotype based vectors, as well as other viral vectors. The rAAV vectors of the invention are particularly advantageous in rAAV readministration and repeat gene therapy.

These and other embodiments and advantages of the invention are described in more detail below. As used throughout this specification and the claims, the term "comprising" is inclusive of other components, elements, integers, steps and the like.

### I. AAV-1 NUCLEIC ACID AND PROTEIN SEQUENCES

The AAV-1 nucleic acid sequences of the invention include the DNA sequences of SEQ ID NO: 1 (Figs. 1A-1C), which consists of 4718 nucleotides. The AAV-1 nucleic acid sequences of the invention further encompass the strand which is complementary to SEQ ID NO: 1, as well as the RNA and cDNA sequences corresponding to SEQ ID NO: 1 and its complementary strand. Also included in the nucleic acid sequences of the invention are natural variants and engineered modifications of SEQ ID NO: 1 and its complementary strand. Such modifications include, for example, labels which are known in the art, methylation, and substitution of one or more of the naturally occurring nucleotides with an analog.

Further included in this invention are nucleic acid sequences which are greater than 85%, preferably at least about 90%, more preferably at least about 95%, and most preferably at least about 98 - 99% identical or homologous to SEQ ID NO:1. The term "percent sequence identity" or "identical" in the context of nucleic acid sequences refers to the residues in the two sequences which are the same when aligned for maximum correspondence. The length of sequence identity comparison may be over the full-length sequence, or a fragment at least about nine nucleotides, usually at least about 20 - 24 nucleotides, at least about 28 - 32 nucleotides, and preferably at least about 36 or more nucleotides. There are a number of different

PCT/US99/25694

5

10

15

20

25

30

algorithms known in the art which can be used to measure nucleotide sequence identity. For instance, polynucleotide sequences can be compared using Fasta, a program in GCG Version 6.1. Fasta provides alignments and percent sequence identity of the regions of the best overlap between the query and search sequences (Pearson, 1990, herein incorporated by reference). For instance, percent sequence identity between nucleic acid sequences can be determined using Fasta with its default parameters (a word size of 6 and the NOPAM factor for the scoring matrix) as provided in GCG Version 6.1, herein incorporated by reference.

The term "substantial homology" or "substantial similarity," when referring to a nucleic acid or fragment thereof, indicates that, when optimally aligned with appropriate nucleotide insertions or deletions with another nucleic acid (or its complementary strand), there is nucleotide sequence identity in at least about 95 - 99% of the sequence.

Also included within the invention are fragments of SEQ ID NO: 1, its complementary strand, cDNA and RNA complementary thereto. Suitable fragments are at least 15 nucleotides in length, and encompass functional fragments which are of biological interest. Certain of these fragments may be identified by reference to Figs. 1A-1C. Examples of particularly desirable functional fragments include the AAV-1 inverted terminal repeat (ITR) sequences of the invention. In contrast to the 145 nt ITRs of AAV-2, AAV-3, and AAV-4, the AAV-1 ITRs have been found to consist of only 143 nucleotides, yet advantageously are characterized by the T-shaped hairpin structure which is believed to be responsible for the ability of the AAV-2 ITRs to direct site-specific integration. In addition, AAV-1 is unique among other AAV serotypes, in that the 5' and 3' ITRs are identical. The full-length 5' ITR sequences of AAV-1 are provided at nucleotides I-143 of SEQ ID NO: 1 (Fig. 1A) and the fulllength 3' ITR sequences of AAV-1 are provided at nt 4576-4718 of SEQ ID NO: 1 (Fig. 1C). One of skill in the art can readily utilize less than the full-length 5' and/or 3' ITR sequences for various purposes and may construct modified ITRs using conventional techniques, e.g., as described for AAV-2 ITRs in Samulski et al, Cell, <u>33</u>:135-143 (1983).

10

15

20

25

Another desirable functional fragment of the AAV-1 genome is the P5 promoter of AAV-1 which has sequences unique among AAV P5 promoters, while maintaining critical regulatory elements and functions. This promoter is located within nt 236 - 299 of SEQ ID NO: 1 (Fig. 1A). Other examples of functional fragments of interest include the sequences at the junction of the rep/cap, e.g., the sequences spanning nt 2306-2223, as well as larger fragments which encompass this junction which may comprise 50 nucleotides on either side of this junction. Still other examples of functional fragments include the sequences encoding the rep proteins. Rep 78 is located in the region of nt 334 - 2306 of SEQ ID NO: 1; Rep 68 is located in the region of nt 334-2272, and contains an intron spanning nt 1924-2220 of SEQ ID NO: 1. Rep 52 is located in the region of nt 1007 - 2304 of SEQ ID NO: 1; rep 40 is located in the region of nt 1007 - 2272, and contains an intron spanning nt 1924-2246 of SEQ ID NO: 1. Also of interest are the sequences encoding the capsid proteins. VP 1 (nt 2223-4431 of SEQ ID NO: 1), VP2 (nt 2634-4432 of SEQ ID NO: 1] and VP3 [nt 2829-4432 of SEQ ID NO: 1]. Other fragments of interest may include the AAV-1 P19 sequences, AAV-1 P40 sequences, the rep binding site, and the terminal resolute site (TRS).

encoded by the AAV-1 nucleic acids of the invention. Particularly desirable proteins include the rep and cap proteins, which are encoded by the nucleotide sequences identified above. These proteins include rep 78 [SEQ ID NO:5], rep 68 [SEQ ID NO:7], rep 52 [SEQ ID NO:9], rep 40 [SEQ ID NO: 11], vpl [SEQ ID NO: 13], vp2 [SEQ ID NO: 15], and vp3 [SEQ IID NO: 17] and functional fragments thereof while the sequences of the rep and cap proteins have been found to be closely related to those of AAV-6, there are differences in the amino acid sequences (see Table 1 below), as well as differences in the recognition of these proteins by the immune system. However, one of skill in the art may readily select other suitable proteins or protein fragments of biological interest. Suitably, such fragments are at least 8 amino acids in length. However, fragments of other desired lengths may be readily utilized.

PCT/US99/25694

5

10

15

20

25

Such fragments may be produced recombinantly or by other suitable means, e.g., chemical synthesis.

The sequences, proteins, and fragments of the invention may be produced by any suitable means, including recombinant production, chemical synthesis, or other synthetic means. Such production methods are within the knowledge of those of skill in the art and are not a limitation of the present invention.

### II. VIRAL VECTORS

In another aspect, the present invention provides vectors which utilize the AAV-1 sequences of the invention, including fragments thereof, for delivery of a heterologous gene or other nucleic acid sequences to a target cell. Suitably, these heterologous sequences (i.e., a transgene) encode a protein or gene product which is capable of being expressed in the target cell. Such a transgene may be constructed in the form of a "minigene". Such a "minigene" includes selected heterologous gene sequences and the other regulatory elements necessary to transcribe the gene and express the gene product in a host cell. Thus, the gene sequences are operatively linked to regulatory components in a manner which permit their transcription. Such components include conventional regulatory elements necessary to drive expression of the transgene in a cell containing the viral vector. The minigene may also contain a selected promoter which is linked to the transgene and located, with other regulatory elements, within the selected viral sequences of the recombinant vector.

Selection of the promoter is a routine matter and is not a limitation of this invention. Useful promoters may be constitutive promoters or regulated (inducible) promoters, which will enable control of the timing and amount of the transgene to be expressed. For example, desirable promoters include the cytomegalovirus (CMV) immediate early promoter/enhancer [see, e.g., Boshart et al, Cell, 41:521-530 (1985)], the Rous sarcoma virus LTR promoter/enhancer, and the chicken cytoplasmic β-actin promoter [T. A. Kost et al, Nucl, Acids Res., 11(23):8287 (1983)]. Still other desirable promoters are the albumin promoter and an AAV P5 promoter. Optionally, the selected promoter is used in conjunction with a heterologous enhancer, e.g., the β-

PCT/US99/25694

5

10

15

20

25

30

actin promoter may be used in conjunction with the CMV enhancer. Yet other suitable or desirable promoters and enhancers may be selected by one of skill in the art.

The minigene may also desirably contain nucleic acid sequences heterologous to the viral vector sequences including sequences providing signals required for efficient polyadenylation of the transcript (poly-A or pA) and introns with functional splice donor and acceptor sites. A common poly-A sequence which is employed in the exemplary vectors of this invention is that derived from the papovavirus SV-40. The poly-A sequence generally is inserted in the minigene downstream of the transgene sequences and upstream of the viral vector sequences. A common intron sequence is also derived from SV-40, and is referred to as the SV40 T intron sequence. A minigene of the present invention may also contain such an intron, desirably located between the promoter/enhancer sequence and the transgene. Selection of these and other common vector elements are conventional [see, e.g., Sambrook et al, "Molecular Cloning. A Laboratory Manual", 2d edit., Cold Spring Harbor Laboratory, New York (1989) and references cited therein] and many such sequences are available from commercial and industrial sources as well as from Genebank.

The selection of the transgene is not a limitation of the present invention. Suitable transgenes may be readily selected from among desirable reporter genes, therapeutic genes, and optionally, genes encoding immunogenic polypeptides. Examples of suitable reporter genes include β-galactosidase (β-gal), an alkaline phosphatase gene, and green fluorescent protein (GFP). Examples of therapeutic genes include, cytokines, growth factors, hormones, and differentiation factors, among others. The transgene may be readily selected by one of skill in the art. See, e.g., WO 98/09657, which identifies other suitable transgenes.

Suitably, the vectors of the invention contain, at a minimum, cassettes which consist of fragments of the AAV-1 sequences and proteins. In one embodiment, a vector of the invention comprises a selected transgene, which is flanked by a 5' ITR and a 3' ITR, at least one of which is an AAV-1 ITR of the invention. Suitably,

10

15

20

25

vectors of the invention may contain a AAV-1 P5 promoter of the invention. In yet another embodiment, a plasmid or vector of the invention contains AAV-1 rep sequences. In still another embodiment, a plasmid or vector of the invention contains at least one of the AAV-1 cap proteins of the invention. Most suitably, these AAV-1-derived vectors are assembled into viral vectors, as described herein.

### A. AAV Viral Vectors

In one aspect, the present invention provides a recombinant AAV-1 viral vector produced using the AAV-1 capsid proteins of the invention. The packaged rAAV-1 virions of the invention may contain, in addition to a selected minigene, other AAV-1 sequences, or may contain sequences from other AAV serotypes.

Methods of generating rAAV virions are well known and the selection of a suitable method is not a limitation on the present invention. See, e.g., K. Fisher et al, <u>J. Virol.</u>, <u>70</u>:520-532 (1993) and US Patent 5,478,745. In one suitable method, a selected host cell is provided with the AAV sequence encoding a rep protein, the gene encoding the AAV cap protein and with the sequences for packaging and subsequent delivery. Desirably, the method utilizes the sequences encoding the AAV-1 rep and/or cap proteins of the invention.

In one embodiment, the rep/cap genes and the sequences for delivery are supplied by co-transfection of vectors carrying these genes and sequences. In one currently preferred embodiment, a cis (vector) plasmid, a trans plasmid containing the rep and cap genes, and a plasmid containing the adenovirus helper genes are co-transfected into a suitable cell line, e.g., 293. Alternatively, one or more of these functions may be provided in trans via separate vectors, or may be found in a suitably engineered packaging cell line.

An exemplary cis plasmid will contain, in 5' to 3' order, AAV 5' ITR, the selected transgene, and AAV 3' ITR. In one desirable embodiment, at least one of the AAV ITRs is a 143 nt AAV-1 ITR. However, other AAV serotype ITRs may be readily selected. Suitably, the full-length ITRs are utilized. However, one of skill in

PCT/US99/25694

5

10

15

20

25

30

the art can readily prepare modified AAV ITRs using conventional techniques.

Similarly, methods for construction of such plasmids is well known to those of skill in the art.

A trans plasmid for use in the production of the rAAV-1 virion particle may be prepared according to known techniques. In one desired embodiment, this plasmid contains the rep and cap proteins of AAV-1, or functional fragments thereof. Alternatively, the rep sequences may be from another selected AAV serotype.

The cis and trans plasmid may then be co-transfected with a wild-type helper virus (e.g., Ad2, Ad5, or a herpesvirus), or more desirably, a replication - defective adenovirus, into a selected host cell. Alternatively, the cis and trans plasmid may be co-transfected into a selected host cell together with a transfected plasmid which provides the necessary helper functions. Selection of a suitable host cell is well within the skill of those in the art and include such mammalian cells as 293 cells, HeLa cells, among others.

Alternatively, the cis plasmid and, optionally the trans plasmid, may be transfected into a packaging cell line which provides the remaining helper functions necessary for production of a rAAV containing the desired AAV-1 sequences of the invention. An example of a suitable packaging cell line, where an AAV-2 capsid is desired, is B-50, which stably expresses AAV-2 rep and cap genes under the control of a homologous P5 promoter. This cell line is characterized by integration into the cellular chromosome of multiple copies (at least 5 copies) of P5-rep-cap gene cassettes in a concatomer form. This B-50 cell line was deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209, on September 18, 1997 under Accession No. CRL-12401 pursuant to the provisions of the Budapest Treaty. However, the present invention is not limited as to the selection of the packaging cell line.

Exemplary transducing vectors based on AAV-1 capsid proteins have been tested both *in vivo and in vitro*, as described in more detail in Example 4. In these studies, it was demonstrated that recombinant AAV vector with an AAV-1 virion can transduce both mouse liver and muscle. These, and other AAV-1 based

gene therapy vectors which may be generated by one of skill in the art are beneficial for gene delivery to selected host cells and gene therapy patients since the neutralization antibodies of AAV-1 present in much of the human population exhibit different patterns from other AAV serotypes and therefore do not neutralize the AAV-1 virions. One of skill in the art may readily prepare other rAAV viral vectors containing the AAV-1 capsid proteins provided herein using a variety of techniques known to those of skill in the art. One may similarly prepare still other rAAV viral vectors containing AAV-1 sequence and AAV capsids of another serotype.

### B. Other Viral Vectors

10

5

One of skill in the art will readily understand that the AAV-1 sequences of the invention can be readily adapted for use in these and other viral vector systems for *in vitro*, *ex vivo or in vivo* gene delivery. Particularly well suited for use in such viral vector systems are the AAV-1 ITR sequences, the AAV-1 rep, the AAV-1 cap, and the AAV-1 P5 promoter sequences.

15

20

For example, in one desirable embodiment, the AAV-1 ITR sequences of the invention may be used in an expression cassette which includes AAV-1 5' ITR, a non-AAV DNA sequences of interest (e.g., a minigene), and 3' ITR and which lacks functional rep/cap. Such a cassette containing an AAV-1 ITR may be located on a plasmid for subsequent transfection into a desired host cell, such as the cis plasmid described above. This expression cassette may further be provided with an AAV capsid of a selected serotype to permit infection of a cell or stably transfected into a desired host cell for packaging of rAAV virions. Such an expression cassette may be readily adapted for use in other viral systems, including adenovirus systems and lentivirus systems. Methods of producing Ad/AAV vectors are well known to those of skill in the art. One desirable method is described in PCT/US95/14018. However, the present invention is not limited to any particular method.

25

30

Another aspect of the present invention is the novel AAV-1 P5 promoter sequences which are located in the region spanning nt 236 - 299 of SEQ ID NO: 1. This promoter is useful in a variety of viral vectors for driving expression of a desired transgene.

PCT/US99/25694

5

10

15

20

25

Similarly, one of skill in the art can readily select other fragments of the AAV-1 genome of the invention for use in a variety of vector systems. Such vectors systems may include, e.g., lentiviruses, retroviruses, poxviruses, vaccinia viruses, and adenoviral systems, among others. Selection of these vector systems is not a limitation of the present invention.

## C. Host Cells And Packaging Cell Lines

In yet another aspect, the present invention provides host cells which may be transiently transfected with AAV-1 nucleic acid sequences of the invention to permit expression of a desired transgene or production of a rAAV particle. For example, a selected host cell may be transfected with the AAV-1 P5 promoter sequences and/or the AAV-1 5' ITR sequences using conventional techniques. Providing AAV helper functions to the transfected cell lines of the invention results in packaging of the rAAV as infectious rAAV particles. Such cell lines may be produced in accordance with known techniques [see, e.g, US Patent No. 5,658,785], making use of the AAV-1 sequences of the invention.

Alternatively, host cells of the invention may be stably transfected with a rAAV expression cassette of the invention, and with copies of AAV-1 rep and cap genes. Suitable parental cell lines include mammalian cell lines and it may be desirable to select host cells from among non-simian mammalian cells. Examples of suitable parental cell lines include, without limitation, HeLa [ATCC CCL 2], A549 [ATCC Accession No. CCL 185], KB [CCL 17], Detroit [e.g., Detroit 510, CCL 72] and WI-38 [CCL 75] cells. These cell lines are all available from the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 USA. Other suitable parent cell lines may be obtained from other sources and may be used to construct stable cell lines containing the P5 and/or AAV rep and cap sequences of the invention.

Recombinant vectors generated as described above are useful for delivery of the DNA of interest to cells.

10

15

20

25

# III. METHODS OF DELIVERING GENES VIA AAV-1 DERIVED VECTORS

In another aspect, the present invention provides a method for delivery of a transgene to a host which involves transfecting or infecting a selected host cell with a recombinant viral vector generated with the AAV-1 sequences (or functional fragments thereof) of the invention. Methods for delivery are well known to those of skill in the art and are not a limitation of the present invention.

In one desirable embodiment, the invention provides a method for AAV-mediated delivery of a transgene to a host. This method involves transfecting or
infecting a selected host cell with a recombinant viral vector containing a selected
transgene under the control of sequences which direct expression thereof and AAV-1
capsid proteins.

Optionally, a sample from the host may be first assayed for the presence of antibodies to a selected AAV serotype. A variety of assay formats for detecting neutralizing antibodies are well known to those of skill in the art. The selection of such an assay is not a limitation of the present invention. See, e.g., Fisher et al, Nature Med., 3(3):306-312 (March 1997) and W. C. Manning et al, Human Gene Therapy, 9:477-485 (March 1, 1998). The results of this assay may be used to determine which AAV vector containing capsid proteins of a particular serotype are preferred for delivery, e.g., by the absence of neutralizing antibodies specific for that capsid serotype.

In one aspect of this method, the delivery of vector with AAV-1 capsid proteins may precede or follow delivery of a gene via a vector with a different serotype AAV capsid protein. Thus, gene delivery via rAAV vectors may be used for repeat gene delivery to a selected host cell. Desirably, subsequently administered rAAV vectors carry the same transgene as the first rAAV vector, but the subsequently administered vectors contain capsid proteins of serotypes which differ from the first vector. For example, if a first vector has AAV-2 capsid proteins, subsequently administered vectors may have capsid proteins selected from among the other serotypes, including AAV-1, AAV-3A, AAV-3B, AAV-4 and AAV-6.

10

15

20

25

30

Thus, a rAAV-1-derived recombinant viral vector of the invention provides an efficient gene transfer vehicle which can deliver a selected transgene to a selected host cell in vivo or ex vivo even where the organism has neutralizing antibodies to one or more AAV serotypes. These compositions are particularly well suited to gene delivery for therapeutic purposes. However, the compositions of the invention may also be useful in immunization. Further, the compositions of the invention may also be used for production of a desired gene product in vitro.

The above-described recombinant vectors may be delivered to host cells according to published methods. An AAV viral vector bearing the selected transgene may be administered to a patient, preferably suspended in a biologically compatible solution or pharmaceutically acceptable delivery vehicle. A suitable vehicle includes sterile saline. Other aqueous and non-aqueous isotonic sterile injection solutions and aqueous and non-aqueous sterile suspensions known to be pharmaceutically acceptable carriers and well known to those of skill in the art may be employed for this purpose.

The viral vectors are administered in sufficient amounts to transfect the cells and to provide sufficient levels of gene transfer and expression to provide a therapeutic benefit without undue adverse effects, or with medically acceptable physiological effects, which can be determined by those skilled in the medical arts. Conventional and pharmaceutically acceptable routes of administration include, but are not limited to, direct delivery to the liver, oral, intranasal, intravenous, intramuscular, subcutaneous, intradermal, and other parental routes of administration. Routes of administration may be combined, if desired.

Dosages of the viral vector will depend primarily on factors such as the condition being treated, the age, weight and health of the patient, and may thus vary among patients. For example, a therapeutically effective human dosage of the viral vector is generally in the range of from about 1 ml to about 100 ml of solution containing concentrations of from about 1 x 10<sup>9</sup> to 1 x 10<sup>16</sup> genomes virus vector. A preferred human dosage may be about 1 x 10<sup>13</sup> to 1 x 10<sup>16</sup> AAV genomes. The dosage will be adjusted to balance the therapeutic benefit against any side effects and

10

15

20

25

such dosages may vary depending upon the therapeutic application for which the recombinant vector is employed. The levels of expression of the transgene can be monitored to determine the frequency of dosage resulting in viral vectors, preferably AAV vectors containing the minigene. Optionally, dosage regimens similar to those described for therapeutic purposes may be utilized for immunization using the compositions of the invention. For *in vitro* production, a desired protein may be obtained from a desired culture following transfection of host cells with a rAAV containing the gene encoding the desired protein and culturing the cell culture under conditions which permits expression. The expressed protein may then be purified and isolated, as desired. Suitable techniques for transfection, cell culturing, purification, and isolation are known to those of skill in the art.

The following examples illustrate several aspects and embodiments of the invention.

# Example 1 - Generation of Infectious Clone of AAV-1

The replicated form DNA of AAV-1 was extracted from 293 cells that were infected by AAV-1 and wild type adenovirus type 5.

### A. Cell Culture and Virus

AAV-free 293 cells and 84-31 cells were provided by the human application laboratory of the University of Pennsylvania. These cells were cultured in Dulbecco's Modified Eagle Medium with 10% fetal bovine serum (Hyclone), penicillin (100 U/ml) and streptomycin at 37°C in a moisturized environment supplied with 5% C0<sub>2</sub>. The 84-31 cell line constitutively expresses adenovirus genes E1a, Elb, E4/ORF6, and has been described previously [K. J. Fisher, J. Virol., 70:520-532 (1996)]. AAV-1 (ATCC VR-645) seed stock was purchased from American Type Culture Collection (ATCC, Manassas, VA). AAV viruses were propagated in 293 cells with wild type Ad5 as a helper virus.

# B. Recombinant AAV Generation

The recombinant AAV viruses were generated by transfection using an adenovirus free method. Briefly, the cis plasmid (with AAV ITR), trans plasmid (with

10

15

20

25

30

AAV rep gene and cap gene) and helper plasmid (pFa13, with essential regions from the adenovirus genome) were simultaneously co-transfected into 293 cells in a ratio of 1:1:2 by calcium phosphate precipitation. The pFa13 helper plasmid has an 8 kb deletion in the adenovirus E2B region and has deletions in most of the late genes. This helper plasmid was generated by deleting the RsrII fragment from pFG140 (Microbix, Canada). Typically, 50 µg of DNA (cis:trans:PFa13 at ratios of 1:1:2, respectively) was transfected onto a 15 cm tissue culture dish. The cells were harvested 96 hours post-transfection, sonicated and treated with 0.5% sodium deoxycholate (37°C for 10 min). Cell lysates were then subjected to two rounds of a CsCl gradient. Peak fractions containing AAV vector were collected, pooled, and dialyzed against PBS before injecting into animals. To make rAAV virus with AAV-1 virion, the pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide rep and cap function.

For the generation of rAAV based on AAV-2, p5E18 was used as the trans plasmid since it greatly improved the rAAV yield. This plasmid, p5E18(2/2), expresses AAV-2 Rep and Cap and contains a P5 promoter relocated to a position 3' to the Cap gene, thereby minimizing expression of Rep78 and Rep68. The strategy was initially described by Li et al, J. Virol., 71:5236-5243 (1997). P5E18(2/2) was constructed in the following way. The previously described pMMTV-trans vector (i.e., the mouse mammary tumor virus promoter substituted for the P5 promoter in an AAV-2-based vector) was digested with SmaI and ClaI, filled in with the Klenow enzyme, and then recircularized with DNA ligase. The resulting construct was digested with XbaI, filled in, and ligated to the blunt-ended BamHI-XbaI fragment from pCR-p5, constructed in the following way. The P5 promoter of AAV was amplified by PCR and the amplified fragment was subsequently cloned into pCR2.1 (Invitrogen) to yield pCR-P5. The helper plasmid pAV1H was constructed by cloning the BfaI fragment of pAAV-2 into pBluescript II-SK(+) at the BcorV and SmaI sites. The 3.0-kb XbaI-KpnI fragment from p5E18(2/2), the 2.3-kb XbaI-KpnI fragment from pAV1H, and the 1.7-kb KpnI fragment from p5E18(2/2) were incorporated into a separate plasmid P5E18(2/1), which contains AAV-2 Rep, AAV-1 Cap, and the

10

15

20

25

30

AAV-2 P5 promoter located 3' to the Cap gene. Plasmid p5E18(2/1) produced 10- to 20-fold higher quantities of the vector than pAV1H (i.e., 10<sup>12</sup> genomes/50 15-cm<sup>2</sup> plates).

### C. <u>DNA Techniques</u>

Hirt DNA extraction was performed as described in the art with minor modification [R.J. Samulski et al., Cell, 33:135-143 (1983)]. More particularly, Hirst solution without SDS was used instead of using original Hirt solution containing SDS. The amount of SDS present in the original Hirst solution was added after the cells had been fully suspended. To construct AAV-1 infectious clone, the Hirt DNA from AAV-1 infected 293 cells was repaired with Klenow enzyme (New England Biolabs) to ensure the ends were blunt. The treated AAV-1 Hirt DNA was then digested with BamHI and cloned into three vectors, respectively. The internal BamHI was cloned into pBlueScript II-SK+ cut with BamHI to get pAV1-BM. The left and right fragments were cloned into pBlueScript II-SK+ cut with BamHI + EcoRV to obtain pAV1-BL and pAV1-BR, respectively. The AAV sequence in these three plasmids were subsequently assembled into the same vector to get AAV-1 infectious clone pAAV-1. The helper plasmid for recombinant AAV-1 virus generation was constructed by cloning the Bfa I fragment of pAAV-1 into pBlueScript II-SK+ at the EcoRV site.

Analysis of the Hirt DNA revealed three bands, a dimer at 9.4 kb, a monomer at 4.7 kb and single-stranded DNA at 1.7 kb, which correlated to different replication forms of AAV-1. The monomer band was excised from the gel and then digested with *BamH*I. This resulted in three fragments of 1.1 kb, 0.8 kb and 2.8 kb. This pattern is in accordance with the description by Bantel-schaal and zur Hausen, Virol., 134(1):52-63 (1984). The 1.1 kb and 2.8 kb *BamH*I fragments were cloned into pBlueScript-KS(+) at *BamH*I and EcoRV site. The internal 0.8 kb fragment was cloned into *BamH*I site of pBlueScript-KS(+).

These three fragments were then subcloned into the same construct to obtain a plasmid (pAAV-1) that contained the full sequence of AAV-1. The pAAV-1 was then tested for its ability to rescue from the plasmid backbone and package

10

15

20

25

infectious virus. The pAAV-1 was then transfected to 293 cells and supplied with adenovirus type as helper at MOI 10. The virus supernatant was used to reinfect 293 cells.

For Southern blot analysis, Hirt DNA was digested with *DpnI* to remove bacteria-borne plasmid and probed with internal *BamHI* fragment of AAV-1. The membrane was then washed at high stringency conditions, which included: twice 30 minutes with 2X SSC, 0.1% SDS at 65°C and twice 30 minutes with 0.1X SSC, 0.1% SDS at 65°C. The membrane was then analyzed by both phosphor image and X-ray autoradiography. The results confirmed that pAAV-1 is indeed an infectious clone of AAV serotype 1.

# Example 2 - Sequencing Analysis of AAV-1

The entire AAV-1 genome was then determined by automatic sequencing and was found to be 4718 nucleotides in length (Figs. 1A-1C). For sequencing, an ABI 373 automatic sequencer as used to determine the sequences for all plasmids and PCR fragments related to this study using the FS dye chemistry. All sequences were confirmed by sequencing both plus and minus strands. These sequences were also confirmed by sequencing two independent clones of pAV-BM, pAV-BL and pAV-BR. Since the replicated form of AAV-1 DNA served as the template for sequence determination, these sequences were also confirmed by sequencing a series of PCR products using original AAV-1 seed stock as a template.

The length of AAV-1 was found to be within the range of the other serotypes: AAV-3 (4726 nucleotides), AAV-4 (4774 nucleotides), AAV-2 (4681 nucleotides), and AAV-6 (4683 nucleotides).

The AAV-1 genome exhibited similarities to other serotypes of adenoassociated viruses. Overall, it shares more than 80% identity with other known AAV
viruses as determined by the computer program Megalign using default settings
[DNASTAR, Madison, WI]. The key features in AAV-2 can also be found in AAV1. First, AAV-1 has the same type of inverted terminal repeat which is capable of
forming T-shaped hairpin structures, despite the differences at the nucleotide level

10

15

20

25

30

(Figs. 2 and 3). The sequences of right ITRs and left ITRs of AAV-1 are identical. The AAV TR sequence is subdivided into A, A', B, B', C, C', D and D' [Bern, cited above].

These AAV ITR sequences are also virtually the same as those found in AAV-6 right ITR, there being one nucleotide difference in each of A and A' sequence, and the last nucleotide of the D sequence. Second, the AAV-2 rep binding motif [GCTCGCTCGCTCGCTG (SEQ ID NO: 20)] is well conserved. Such motif can also be found in the human chromosome 19 AAV-2 pre-integration region. Finally, non-structural and structural coding regions, and regulatory elements similar to those of other AAV serotypes also exist in AAV-1 genome.

Although the overall features of AAV terminal repeats are very much conserved, the total length of the AAV terminal repeat exhibits divergence. The terminal repeat of AAV-1 consists of 143 nucleotides while those of AAV-2, AAV-3, and AAV-4 are about 145 or 146 nucleotides. The loop region of AAV-1 ITR most closely resembles that of AAV-4 in that it also uses TCT instead of the TTT found in AAV-2 and AAV-3. The possibility of sequencing error was eliminated using restriction enzyme digestion, since these three nucleotides are part of the SacI site (gagete; nt 69-74 of SEQ ID NO: 1). The p5 promoter region of AAV-1 shows more variations in nucleotide sequences with other AAV serotypes. However, it still maintains the critical regulatory elements. The two copies of YY1 [See, Fig. 1A-1C] sites seemed to be preserved in all known AAV serotypes, which have been shown to be involved in regulating AAV gene expression. In AAV-4, there are 56 additional nucleotides inserted between YY1 and E-box/USF site, while in AAV-1, there are 26 additional nucleotides inserted before the E-box/USF site. The p19 promoter, p40 promoter and polyA can also be identified from the AAV-1 genome by analogy to known AAV serotypes, which are also highly conserved.

Thus, the analysis of AAV terminal repeats of various serotypes showed that the A and A' sequence is very much conserved. One of the reasons may be the Rep binding motif (GCTC)<sub>3</sub>GCTG [SEQ ID NO: 20]. These sequences appear to be essential for AAV DNA replication and site-specific integration. The same sequence

15

20

25

has also been shown to be preserved in a monkey genome [Samulski, personal communication]. The first 8 nucleotides of the D sequence are also identical in all known AAV serotypes. This is in accordance with the observation of the Srivastava group that only the first 10 nucleotides are essential for AAV packaging [X.S. Wang et al, J. Virol., 71:3077-3082 (1997); X.S. Wang et al, J. Virol., 71:1140-1146 (1997)]. The function of the rest of the D sequences still remain unclear. They may be somehow related to their tissue specificities. The variation of nucleotide in B and C sequence may also suggest that the secondary structure of the ITRs is more critical for its biological function, which has been demonstrated in many previous publications.

### Example 3 - Comparison of AAV-1 Sequences

The nucleotide sequences of AAV-1, obtained as described above, were compared with known AAV sequences, including AAV-2, AAV-4 and AAV-6 using DNA Star Megalign. This comparison revealed a stretch of 71 identical nucleotides shared by AAV-1, AAV-2 and AAV-6. See, Figs. 1A-1C.

This comparison further suggested that AAV-6 is a hybrid formed by homologous recombination of AAV-1 and AAV-2. See, Figs. 3A and 3B. These nucleotides divide the AAV-6 genome into two regions. The 5' half of AAV-6 of 522 nucleotides is identical to that of AAV-2 except in 2 positions. The 3' half of AAV-6 including the majority of the rep gene, complete cap gene and 3' ITR is 98% identical to AAV-1.

Biologically, such recombination may enable AAV-1 to acquire the ability to transmit through the human population. It is also interesting to note that the ITRs of AAV-6 comprise one AAV-1 ITR and one AAV-2 ITR. The replication model of defective parvovirus can maintain this special arrangement. Studies on AAV integration have shown that a majority of AAV integrants carries deletions in at least one of the terminal repeats. These deletions have been shown to be able to be repaired through gene conversion using the other intact terminal repeat as a template. Therefore, it would be very difficult to maintain AAV-6 as a homogenous population

when an integrated copy of AAV-6 is rescued from host cells with helper virus infection. The AAV-6 with two identical AAV-2 ITRs or two identical AAV-1 ITRs should be the dominant variants. The AAV-6 with two AAV-1 ITRs has been observed by Russell's group [Rutledge, cited above (1998)]. So far there is no report on AAV-6 with two AAV-2 ITRs. Acquirement of AAV-2 P5 promoter by AAV-6 may have explained that AAV-6 have been isolated from human origin while AAV-1 with the same virion has not. The regulation of P5 promoter between different species of AAV may be different *in vivo*. This observation suggests the capsid proteins of AAV were not the only determinants for tissue specificity.

10

15

5

Although it is clear that AAV-6 is a hybrid of AAV-1 and AAV-2, AAV-6 has already exhibited divergence from either AAV-1 or AAV-2. There are two nucleotide differences between AAV-6 and AAV-2 in their first 450 nucleotides. There are about 1% differences between AAV-6 and AAV-1 in nucleotide levels from nucleotides 522 to the 3' end. There also exists a quite divergent region (nucleotide 4486-4593) between AAV-6 and AAV-1 (Figs. 1A-1C). This region does not encode any known proteins for AAVs. These differences in nucleotide sequences may suggest that AAV-6 and AAV-1 have gone through some evolution since the recombination took place. Another possible explanation is that there exists another variant of AAV-1 which has yet to be identified. So far, there is no evidence to rule out either possibility. It is still unknown if other hybrids (AAV-2 to AAV-4, etc.) existed in nature.

20

The coding region of AAV-1 was deduced by comparison with other known AAV serotypes. Table 1 illustrates the coding region differences between AAV-1 and AAV-6. The amino acid residues are deduced according to AAV-2.

25

With reference to the amino acid position of AAV-1, Table 1 lists the amino acids of AAV-1 which have been changed to the corresponding ones of AAV-6. The amino acids of AAV-1 are shown to the left of the arrow. Reference may be made to SEQ ID NO: 5 of the amino acid sequence of AAV-1 Rep 78 and to SEQ ID NO: 13 for the amino acid sequence of AAV-1 VP1.

PCT/US99/25694

Table 1

Coding region variations between AAV-1 and AAV-6

24

Rep protein (Rep78)			Cap protein (VP1)		
Position(s)	Amino acids		Position(s)	Amino acids	
28	S→N		129	L→F	
191	Q-H		418	E→D	
192	H-D		531	E→K	
308	E-D		584	F→L	
			598	A-V	
			642	N→H	

It was surprising to see that the sequence of the AAV-1 coding region is
almost identical to that of AAV-6 from position 452 to the end of coding region
(99%). The first 508 nucleotides of AAV-6 have been shown to be identical to those
of AAV-2 [Rutledge, cited above (1998)]. Since the components of AAV-6 genome
seemed to be AAV-2 left ITR - AAV-2 p5 promoter - AAV-1 coding region - AAV1 right ITR, it was concluded that AAV-6 is a naturally occurred hybrid between
AAV-1 and AAV-2.

### Example 4 - Gene Therapy Vector Based on AAV-1

Recombinant gene transfer vectors based on AAV-1 viruses were constructed by the methods described in Example 1. To produce a hybrid recombinant virus with AAV-1 virion and AAV-2 ITR, the AAV-1 trans plasmid (pAV1H) and the AAV-2 cis-lacZ plasmid (with AAV-2 ITR) were used. The AAV-2 ITR was used in this vector in view of its known ability to direct site-specific integration. Also constructed for use in this experiment was an AAV-1 vector carrying the green fluorescent protein (GFP) marker gene under the control of the immediate early promoter of CMV using pAV1H as the trans plasmid.

5

20

10

15

20

25

### A. rAAV-1 Viruses Transfect Host Cells in Vitro

84-31 cells, which are subclones of 293 cells (which express adenovirus E1a, E1b) which stably express E4/ORF5, were infected with rAAV-1 GFP or rAAV-lacZ. High levels of expression of GFP and lacZ was detected in the cultured 84-31 cells. This suggested that rAAV-1 based vector was very similar to AAV-2 based vectors in ability to infect and expression levels.

### B. rAAV-1 Viruses Transfect Cells in Vivo

The performance of AAV-1 based vectors was also tested *in vivo*. The rAAV-1 CMV-α1AT virus was constructed as follows. The EcoRI fragment of pAT85 (ATCC) containing human α1-antitrypsin (α1AT) cDNA fragment was blunted and cloned into PCR (Promega) at a Smal site to obtain PCR-α1AT. The CMV promoter was cloned into PCR-α1AT at the Xbal site. The Alb-α1AT expression cassette was removed by Xhol and ClaI and cloned into pAV1H at the Xbal site. This vector plasmid was used to generate AAV-1-CMV-α1AT virus used in the experiment described below.

For screening human antibodies against AAV, purified AAV virus is lysed with Ripa buffer (10 mM Tris pH 8.2, 1% Triton X-100, 1% SDS, 0.15 M NaCl) and separated in 10% SDS-PAGE gel. The heat inactivated human serum was used at a 1 to 1000 dilution in this assay. The rAAV-1 CMV-α1AT viruses were injected into Rag-1 mice through tail vein injection at different dosages. The concentration of human α1-antitrypsin in mouse serum was measured using ELISA. The coating antibody is rabbit anti-human human α1-antitrypsin (Sigma). The goatantihuman α1-antitrypsin (Sigma) was used as the primary detection antibodies. The sensitivity of this assay is around 0.3 ng/ml to 30 ng/ml. The expression of human α-antitrypsin in mouse blood can be detected in a very encouraging level. This result is shown in Table 2.

10

15

20

25

Table 2
Human Antitrypsin Expressed in Mouse Liver

Amount of virus injected	Week 2 (ng/ml)	Week 4 (ng/ml)
2x10 <sup>10</sup> genomes	214.2	171.4
1x10 <sup>10</sup> genomes	117.8	109.8
5x10 <sup>10</sup> genomes	64.5	67.8
2.5x10 <sup>10</sup> genomes	<b>30</b> .9	58.4

rAAV-1 CMV-lacZ viruses were also injected into the muscle of C57BL6 mice and similar results were obtained. Collectively, these results suggested that AAV-1 based vector would be appropriate for both liver and muscle gene delivery.

# Example 5 - Neutralizing Antibodies Against AAV-1

Simple and quantitative assays for neutralizing antibodies (NAB) to AAV-1 and AAV-2 were developed with recombinant vectors. A total of 33 rhesus monkeys and 77 normal human subjects were screened.

### A. Nonhuman Primates

Wild-caught juvenile rhesus monkeys were purchased from Covance (Alice, Tex.) and LABS of Virginia (Yemassee, SC) and kept in full quarantine. The monkeys weighed approximately 3 to 4 kg. The nonhuman primates used in the Institute for Human Gene Therapy research program are purposefully bred in the United States from specific-pathogen-free closed colonies. All vendors are US Department of Agriculture class A dealers. The rhesus macaques are therefore not infected with important simian pathogens, including the tuberculosis agent, major simian lentiviruses (simian immunodeficiency virus and simian retroviruses), and cercopithecine herpesvirus. The animals are also free of internal and external parasites. The excellent health status of these premium animals minimized the potential for extraneous variables. For this study, serum was obtained from monkeys prior to initiation of any protocol.

10

15

20

25

NAB titers were analyzed by assessing the ability of serum antibody to inhibit the transduction of reporter virus expressing green fluorescent protein (GFP) (AAV1-GFP or AAV2-GFP) into 84-31 cells. Various dilutions of antibodies preincubated with reporter virus for 1 hour at 37°C were added to 90% confluent cell cultures. Cells were incubated for 48 hours and the expression of green fluorescent protein was measured by FluoroImaging (Molecular Dynamics). NAB titers were calculated as the highest dilution at which 50% of the cells stained green.

Analysis of NAB in rhesus monkeys showed that 61% of animals tested positive for AAV-1; a minority (24%) has NAB to AAV-2. Over one-third of animals had antibodies to AAV-1 but not AAV-2 (i.e., were monospecific for AAV-1), whereas no animals were positive for AAV-2 without reacting to AAV-1. These data support the hypothesis that AAV-1 is endemic in rhesus monkeys. The presence of true AAV-2 infections in this group of nonhuman primates is less clear, since cross-neutralizing activity of an AAV-1 response to AAV-2 can not be ruled out. It is interesting that there is a linear relationship between AAV-2 NAB and AAV-1 NAB in animals that had both.

### B. Humans

For these neutralization antibody assays, human serum samples were incubated at 56°C for 30 min to inactivate complement and then diluted in DMEM. The virus (rAAV or rAd with either lacZ or GFP) was then mixed with each serum dilution (20X, 400X, 2000X, 4000X, etc.) and incubated for 1 hour at 37°C before applied to 90% confluent cultures of 84-31 cells (for AAV) or Hela cells (for adenovirus) in 96-well plates. After 60 minutes of incubation at culture condition, 100 µl additional media containing 20% FCS was added to make final culture media containing 10% FCS.

The result is summarized in Table 3.

Table 3

Adenovirus	AAV-1	AAV-2	# of samples	Percentage
_	_	_	41	53.2%
+	-		16	20.8%
-	+	-	0	0.0%
-	-	+	2	2.6%
_	+	+	2	2.6%
+	-	+	3	3.9%
+	+	-	0	0.0%
+	+ '	+	13	16.9%
		Total	77	100%

The human neutralizing antibodies against these three viruses seemed to be unrelated since the existence of neutralizing antibodies against AAV are not indications for antibodies against adenovirus. However, AAV requires adenovirus as helper virus, in most of the cases, the neutralizing antibodies against AAV correlated with the existence of neutralizing antibodies to adenovirus. Among the 77 human serum samples screened, 41% of the samples can neutralize the infectivity of recombinant adenovirus based on Ad5. 15/77 (19%) of serum samples can neutralize the transduction of rAAV-1 while 20/77 (20%) of the samples inhibit rAAV-2 transduction at 1 to 80 dilutions or higher. All serum samples positive in neutralizing antibodies for AAV-1 in are also positive for AAV-2. However, there are five (6%) rAAV-2 positive samples that failed to neutralize rAAV-1. In samples that are positive for neutralizing antibodies, the titer of antibodies also varied in the positive ones. The results from screening human sera for antibodies against AAVs supported the conclusion that AAV-1 presents the same epitome as that of AAV-2 to interact

5

10

15

25

20

10

15

20

25

with cellular receptors since AAV-1 neutralizing human serums can also decrease the infectivity of AAV-2. However, the profile of neutralizing antibodies for these AAVs is not identical, there are additional specific receptors for each AAV serotype.

# Example 6 - Recombinant AAV Viruses Exhibit Tissue Tropism

The recombinant AAV-1 vectors of the invention and the recombinant AAV-2 vectors [containing the gene encoding human α1-antitrypsin (α1AT) or murine erythropoietin (Epo) from a cytomegalovirus-enhanced β-actin promoter (CB)] were evaluated in a direct comparison to equivalent copies of AAV-2 vectors containing the same vector genes.

Recombinant viruses with AAV-1 capsids were constructed using the techniques in Example 1. To make rAAV with AAV-1 virions, pAV1H or p5E18 (2/1) was used as the *trans* plasmid to provide Rep and Cap functions. For the generation of the rAAV based on AAV-2, p5E18(2/2) was used as the *trans* plasmid, since it greatly improved the rAAV yield. [Early experiments indicated similar *in vivo* performances of AAV-1 vectors produced with pAV1H and p5E19 (2/1). All subsequent studies used AAV-1 vectors derived from p5E18(2/1) because of the increased yield.]

Equivalent stocks of the AAV-1 and AAV-2 vectors were injected intramuscularly (5 x 10<sup>10</sup> genomes) or liver via the portal circulation (1 x 10<sup>11</sup> genomes) into immunodeficient mice, and the animals (four groups) were analyzed on day 30 for expression of transgene. See, Figs. 4A and 4B.

AAV-2 vectors consistently produced 10- to 50-fold more serum erythropoietin or α1-antitrypsin when injected into liver compared to muscle. (However, the AAV-1-delivered genes did achieve acceptable expression levels in the liver.) This result was very different from that for AAV-1 vectors, with which muscle expression was equivalent to or greater than liver expression. In fact, AAV-1 outperformed AAV-2 in muscle when equivalent titers based on genomes were administered.

10

15

20

25

30

### Example 7 - Gene Delivery via rAAV-1

C57BL/6 mice (6- to 8-week old males, Jackson Laboratories) were analyzed for AAV mediated gene transfer to liver following intrasplenic injection of vector (i.e., targeted to liver). A total of 10<sup>11</sup> genome equivalents of rAAV-1 or rAAV-2 vector were injected into the circulation in 100 μl buffered saline. The first vector contained either an AAV-1 capsid or an AAV-2 capsid and expressed α1AT under the control of the chicken β-actin (CB) promoter. Day 28 sera were analyzed for antibodies against AAV-1 or AAV-2 and serum α1AT levels were checked. Animals were then injected with an AAV-1 or AAV-2 construct expressing erythropoietin (Epo, also under the control of the CB promoter). One month later sera was analyzed for serum levels of Epo. The following groups were analyzed (Figs. 5A-5D).

In Group 1, vector 1 was AAV-2 expressing a1AT and vector 2 was AAV-2 expressing Epo. Animals generated antibodies against AAV-2 following the first vector administration which prevented the readministration of the AAV-2 based vector. There was no evidence for cross-neutralizing the antibody to AAV-1.

In Group 2, vector 1 was AAV-1 expressing a1AT while vector 2 was AAV-1 expressing Epo. The first vector administration did result in significant a1AT expression at one month associated with antibodies to neutralizing antibodies to AAV-1. The animals were not successfully readministered with the AAV-1 Epo expressing construct.

In Group 3, the effectiveness of an AAV-2 vector expressing Epo injected into a naive animal was measured. The animals were injected with PBS and injected with AAV-2 Epo vector at day 28 and analyzed for Epo expression one month later. The neutralizing antibodies were evaluated at day 28 so we did not expect to see anything since they received PBS with the first vector injection. This shows that in naive animals AAV-2 is very efficient at transferring the Epo gene as demonstrated by high level of serum Epo one month later.

Group 4 was an experiment similar to Group 3 in which the animals originally received PBS for vector 1 and then the AAV-1 expressing Epo construct 28 days later. At the time of vector injection, there obviously were no antibodies to either

10

15

20

AAV-1 or AAV-2. The AAV-1 based vector was capable of generating significant expression of Epo when measured one month later.

Group 5 is a cross-over experiment where the initial vector is AAV-2 expressing α1AT followed by the AAV-1 construct expressing Epo. The animals, as expected, were efficiently infected with the AAV-2 vector expressing α1AT as shown by high levels of the protein in blood at 28 days. This was associated with significant neutralizing antibodies to AAV-2. Importantly, the animals were successfully administered AAV-1 following the AAV-2 vector as shown by the presence of Epo in serum 28 days following the second vector administration. At the time of this vector administration, there was high level AAV-2 neutralizing antibodies and very low cross-reaction to AAV-1. The level of Epo was slightly diminished possibly due to a small amount of cross-reactivity. Group 6 was the opposite cross-over experiment in which the initial vector was AAV-1 based, whereas the second experiment was AAV-2 based. The AAV-1 vector did lead to significant gene expression of α1AT, which also resulted in high level AAV-1 neutralizing antibody. The animals were very efficiently administered AAV-2 following the initial AAV-1 vector as evidenced by high level Epo.

A substantially identical experiment was performed in muscle in which 5 x 10<sup>10</sup> genomes were injected into the tibialis anterior of C57BL/6 mice as a model for muscle directed gene therapy. The results are illustrated in Figs. 6A-6D and are essentially the same as for liver.

In summary, this experiment demonstrates the utility of using an AAV-1 vector in patients who have pre-existing antibodies to AAV-2 or who had initially received an AAV-2 vector and need readministration.

### 25 Example 8 - Construction of Recombinant Viruses Containing AAV-1 ITRs

This example illustrates the construction of recombinant AAV vectors which contain AAV-1 ITRs of the invention.

An AAV-1 cis plasmid is constructed as follows. A 160 bp Xho-NruI AAV-1 fragment containing the AAV-1 5' ITR is obtained from pAV1-BL. pAV1-BL was

10

15

20

generated as described in Example 1. The Xho-NruI fragment is then cloned into a second pAV1-BL plasmid at an XbaI site to provide the plasmid with two AAV-1 ITRs. The desired transgene is then cloned into the modified pAV-1BL at the NruI and BamHI site, which is located between the AAV-1 ITR sequences. The resulting AAV-1 cis plasmid contains AAV-1 ITRs flanking the transgene and lacks functional AAV-1 rep and cap.

Recombinant AAV is produced by simultaneously transfecting three plasmids into 293 cells. These include the AAV-1 cis plasmid described above; a trans plasmid which provides AAV rep/cap functions and lacks AAV ITRs; and a plasmid providing adenovirus helper functions. The rep and/or cap functions may be provided in trans by AAV-1 or another AAV serotype, depending on the immunity profile of the intended recipient. Alternatively, the rep or cap functions may be provided in cis by AAV-1 or another serotype, again depending on the patient's immunity profile.

In a typical cotransfection, 50 µg of DNA (cis:trans:helper at ratios of 1:1:2, respectively) is transfected onto a 15 cm tissue culture dish. Cells are harvested 96 hours post transfection, sonicated and treated with 0.5% sodium deoxycholate (37° for 10 min). Cell lysates are then subjected to 2-3 rounds of ultracentrifugation in a cesium gradient. Peak fractions containing rAAV are collected, pooled and dialyzed against PBS. A typical yield is 1 x 10<sup>13</sup> genomes/10° cells.

Using this method, one recombinant virus construct is prepared which contains the AAV-1 ITRs flanking the transgene, with an AAV-1 capsid. Another recombinant virus construct is prepared with contains the AAV-1 ITRs flanking the transgene, with an AAV-2 capsid.

All publications cited in this specification are incorporated herein by reference.

While the invention has been described with reference to a particularly preferred embodiments, it will be appreciated that modifications can be made without departing from the spirit of the invention. Such modifications are intended to fall within the scope of the claims.

What is claimed is:

- 1. An isolated AAV-1 nucleic acid molecule comprising a sequence selected from the group consisting of:
  - (a) SEQ ID NO: 1;
  - (b) a DNA sequence complementary to SEQ ID NO: 1;
  - (c) cDNA complementary to (a) or (b); and
  - (d) RNA complementary to any of (a) to (c).
- 2. A nucleic acid molecule comprising an AAV-1 inverted terminal repeat (ITR) sequence selected from the group consisting of:
  - (a) nt 1 to 143 of SEQ ID NO: 1;
  - (b) nt 4576 to 4718 of SEQ ID NO: 1;
  - (c) a nucleic acid sequence complementary to (a) or (b); and
  - (d) a functional fragment of (a), (b), or (c).
- 3. A recombinant vector comprising a 5' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said 1TR has the sequence selected from the group consisting of:
  - (a) nt 1 to 143 of SEQ ID NO: 1;
  - (b) a nucleic acid sequence complementary to (a); and
  - (c) a functional fragment of (a) or (b).
- 4. The recombinant vector according to claim 3, wherein said vector further comprises a 3' AAV-1 ITR.

- 5. A recombinant vector comprising a 3' AAV-1 inverted terminal repeat (ITR) and a selected transgene, wherein said ITR has the sequence selected from the group consisting of:
  - (a) nt 4576 to 4718 of SEQ ID NO: 1;
  - (b) a nucleic acid sequence complementary to (a); and
  - (c) a functional fragment of (a) or (b).
- 6. The recombinant vector according to claim 5, wherein said vector further comprises a 5' AAV-1 ITR.
- 7. The recombinant vector according to any of claims 3-6, wherein said vector further comprises AAV-1 capsid proteins having the sequence of SEQ ID NO: 13, 15 or 17 or functional fragments thereof.
- 8. The recombinant vector according to any of claims 3-6, wherein said vector further comprises adenovirus sequences.
- 9. A recombinant vector comprising an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1 or a functional fragment thereof.
- 10. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said cap coding region comprises at least one member is selected from the group consisting of:
  - (a) vp1, nt 2223 to 4431 of SEQ ID NO: 1;
  - (b) vp2, nt 2634 to 4432 of SEQ ID NO: 1; and
  - (c) vp3, nt 2829 to 4432 of SEQ ID NO: 1.

- 11. A nucleic acid molecule encoding AAV-1 helper functions, said molecule comprising an AAV rep coding region and an AAV cap coding region, wherein said rep coding region comprises an AAV-1 rep coding region comprising at least one member selected from the group consisting of:
  - (a) rep 78, nt 335 to 2304 of SEQ ID NO: 1;
- (b) rep 68, nt 335 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto;
  - (c) rep 52, nt 1007 to 2304 of SEQ ID NO: 1; and
- (d) rep 40, nt 1007 to 2272 of SEQ ID NO: 1 or the cDNA corresponding thereto.
- 12. A host cell transduced with a recombinant viral vector according to any of claims 3-6.
- 13. A host cell transduced with a nucleic acid molecule according to any of claims 1, 2, 10 or 11.
- 14. A host cell stably transduced with an AAV-1 P5 promoter having the sequence of nt 236 to 299 of SEQ ID NO: 1.
- 15. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to any of claims 3-6.
- 16. A pharmaceutical composition comprising a carrier and a virus comprising the vector according to claim 7.
- 17. A pharmaceutical-composition comprising a carrier and a virus comprising the vector according to claim 8.

- 18. A method for AAV-mediated delivery of a transgene comprising the step of delivering to a host cell an AAV virion which comprises:
- (a) a capsid comprising at least one capsid protein encoded by an AAV-1 cap gene; and
- (b) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 19. A method for AAV-mediated delivery of a transgene to a host comprising the steps of:
- (a) assaying a sample from the host to determine the presence of neutralizing antibodies specific against any serotype of AAV; and
  - (b) delivering to the host an AAV virion which comprises:
- (i) a capsid comprising at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has no antibodies as determined in step (a); and
- (ii) a DNA molecule comprising a transgene under the control of regulatory sequences directing its expression.
- 20. The method according to claim 19, comprising the additional step of repeating steps (a) and (b).
- 21. Use of an AAV virion which comprises a capsid comprising (a) at least one capsid protein encoded by a cap gene of an AAV serotype against which the host has antibodies, and (b) a DNA molecule comprising a transgene operably linked to regulatory sequences directing its expression,

in the preparation of a medicament for delivery of a transgene to a host, wherein said host has no preexisting neutralizing antibodies against the AAV serotype of said cap gene.

- 22. A method for delivery of a transgene comprising the step of delivering to a host cell a recombinant virus comprising a recombinant vector according to any of claims 3-8.
- A method for producing a selected gene product comprising the steps of transfecting a mammalian cell with the molecule according to claim 1 or a functional fragment thereof and culturing said cell under conditions suitable to express said gene product.

## FIGIA

	Reportation -	
	ttgcccactccctcttgcgcgctcgctcggttggggcctgcggaccaaaggtcgcaaaggtcgcagagctctgctctgccggcccaacqggggggcgcaaaggggggccaaaggtcgcagagagagccaaaggtcgcagagagag	110
AAV-1		128
AAT - 6	.ac.ag.gc	
*****	E hov//ISF	
	TRS  E box/USF  Fyc. backters et backtogget and too control co	
	THE NAME OF THE PROPERTY OF TH	127
AAW. 3	F	222
AAV-6	######################################	
221	Ren7N/68	
	YYI FAIR TIME TO THE TOTAL	156
AAV-1	GACATITIOCENCACCACUTOCCATTTACOCCAGTGAOCGACCACCACCACCACCACCACCACCACCACCACCACCAC	142
AAV-2		341
AAV-6		
	TEATCALOGTOCCCA.OCCIACCTOCIACIAGEA.CCTGCCOGGCATTTCTGA.CTCGTTTGTGIACCTGGGTTGGCCCGAGAAGAATCGGAGCTGCCCCCCGAGTTCTGAATCTGA	476
AAV-L	TEXTELLOGICCCCACCUACCTGCCCGCCCCCCCCCCCTTCCCGTTGCGLACCCCCCACLAGGLACGLACCCACTACCCCCCACAAGGLACGCACTACCCCCCACAAGGLACGCACTACCCCCCACAAGGLACGCACACACACACACACACACACAC	462
AAV-3	TG.A	461
444.	TGG	
	• •	
	TTGAGCAGOCACCCCTCACCGTGGCCCACAACCTGCAGCCCCCACTTCCTTGTTCCAATGGCCCCCCGTGAGTAACGCCCCCCGGAGGCCCCTCTTCTTTGTTCCAGAAGGGCGGAGTCCCAGAAGGCCGAGAGCCCAGAAGCCCCAGAAGCCCAGAAGCCCAGAAGCCCAGAAAGCCCAGAAAGCCCAGAAAACCAGAAAAAGCCCAGAAAAACCAGAAAAAA	594
<b>77A-7</b>	TTGAGCAGOCACCCCTEACCOTGCCCAGAACCTGCAGCGCCACTTCCTGGTCGAACCCCCCCGCTGAGTAAGCCCCCAGAACCTGCAGCACCTGCAACCTGCAGCACCTGCAACCTGCAGCACCTGCAACCTGCAGCACCTGCAGCACCTGCAGCACCTGCAACCTCAACCTGCAACCTGCAACCTGCAACCTGCAACCTGCAACCTGCAACCTGCAACCTGCAACCTGCAACCT	582
YYA-1		582
AAV-6	······································	
	ACTICCACTICCATATICTOGIOGIACCACGOGOGICAAATCCATGGGGCTGCTGGGGGCGGATCGAGGGCGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGGACCATGTACGGGACCATGTACGGGACCATGTACGGGACCATGTACGGGACCATGTACGGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGGACCATGTACGACGACCATGTACGACGACCATGTACGACGACCATGTACGACGACCATGTACGACGACCATGTACGACGACATGTACGACGACATGTACGACGACATGTACGACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACGACATGTACATGTACGACATGTACATGTACGACATGTACATATACATGTACATATACATGTACATATACATGTACATATACATATACATATACATATATACATATATAT	716
AAV-1	ACTICCACTICCATATICTOGROUNGACCACGOOGUTCAAATCATGGTOCTOGCOCGCTTC.TGGGTCAGATTAGATTAGGTCAGATTAGATTAGGTCAGATTAGATTAGATTAGGTCAGATTAGA	702
<b>77A-3</b>	A.GCG.GCACGTTAT	701
74.4		
<b>-</b>	TECCEMETACTICCOCCUMENTALICCOCCUMENTATION CONTRACTOCOCCUMENTALICCOCCUME	836
AAV-1	TECCCLACTECCOCTUACCAMMICCOTTALTOCCCCCACOCOCCACTOCTCACTACTACTACTACTACTACTACTACTACTACTACTA	622
MV-3		121
MA-0	••••••	
	P19/TATA P19 RNA	
	CONTROL TAXABLE PROPERTY AND CONCENTRAL ACCIDENCE AND CONCENTRAL C	730
AAV-1	CGTOTALCAROGROUNGTATATANOCOCCTOTT TO ACCTOROCOCAGOCCAGOCCAGOCCAGOCCAGOCCAGOCCAGOC	243
MY-4	C	7.
MV-0		
	Rep 52/40	3074
14V-1	REDIMO REDIMO RESIDENCIA DE CONTROL DE CONTR	1043
LAV-3	ACCCUATIFICACCOCCUTCATACCCCCATACCCCCATACCCCCCCCCCC	2061
MY-6	.T	
	TALL TOUR TRANSPORT OF THE PROPERTY OF THE PRO	1196
LAV-1	CCTCOTHCATCTCCTTCAACOCCCCTTCCAACTCCCCCACTCACACTCACCCCCCCC	1102
MY-2	CCTCGFRCATCTCCTTCAACCCCTTCCAACTCCGGGTCCGGATCAAGGCCGCTCTGGGCAATCCCGGCAAGATCATGGCCGCAAGATCATGGCCGCAAGATCATGGCCGCAACTCTGGCCAACTCTGCAACTCTAACTCTGCAACTCTGCAACTCT	1181
AV-6		
_		
~	CTCCGCCCCCCCCCATTAAAACCAACCCCATCTTACCCCATCTTGCACCTTACACCCCACCACCTTCCCCCCCC	2316
AV-1	CTCCSCCCCCCCCACATAAAACCAACCCATCTACCCCATC.TGGAGCTGAACCCCTACGAACCTCCTACACACACACACAC	1302
WY-2	AGCOTG.ATCC.GTGTTAAATTAAGTCDATG.CTAA	1301
TA-6	AGCSTG.ATCC.GTGTTANATTAAGTCCNATG.CT	
	ACACCAMETOCCOMO TROCCCOCCACCACCACCACCACCACCACCACCACCCACCC	2436
74-7	ACACCAMIC POCCOCCOCCACCACCACCACCACCACCACCACCACCACCA	1422
<b>7A-5</b>	ACROCANCIDED FIGURE CONTROL OF A CONTROL OF	1421
<b>4-4</b>	***************************************	
	ATTOCOTOCACAAGATOGTCATCTOGTGGGGAGGGGCAAGATCAAGGCCAAGGTCGTGGAGTCGGGGAGGGGGGGG	2556
7A-5	ATTOCOTOCAL ACTION CONTROL ACTION CO	1342
74-1	ATTOCHICALGARGATGGTGGTGGGGGGGGGGGGGGCAGGTGAGGGCGAGGCGAGGCGATGCGGGAGGCGATGCTGGGGGGGG	1541
74-6	***************************************	
	·	
	CCCCCCAGATCGACCCCCCCCCCCCCCCACCCCCACACACA	1676
44 - J	GCCCCCACATCCACCCCCCCCCCCCCCCCCCCCCCCCCC	1663
VA-3	.d	1661
7A-2		
	TTCLACTCACCCCCTCTGGCCATGLCTTTGGCAAGGTCACAAAGCAGGAAAGTCAAGAAGTCTTCCGCTGGCCGCAGATCACGTGACGCAAGTGCCCCAAGTTCTACGTCACA	1796
AV-1	TTCLACTCCCCCCCCTCTCGCCATCLCTTTCGCCACGTCACAAACACCAAGTCAAAGAATTCTTCCCTCGCCCACAACACACAACACAACACAACACAACACAACACAACAC	1702
AV-Z	······································	1781
AV-6		
	P40TATA P40 RNA	
		1916
	AG A T T T C C.G T A AC AC.	1077
V-2	ACOGTOCACCELACAAAAGACCCCCCCGATGACGCCGATAAAAGCGACCCCAACCCCCCCC	1991

## FIG 1B

	CONCROTRACIALACAARTITECTCUTCACGCCCCCCATGCTTCCAGATOCTOTTTCCCTGCAAGACATGCAGAAATGAATCAGAATTTCAACATTTCAACGCACGGAACGAGAAAAAAAA	. 2019
	ACTOTICMANGIOCITECCCCCCCCCTGTCAGAATCTCAACCCCGTCGTCAGAAAGAGGACGTATCCGAAAAGACCTCTGTGCCATTCATCATCATCTCCCGGGGCCGGGCCTGCCCGAGATTGCTTGC	2153
		· JUJ
AAV.		2138
	Rep78 stop VP1 V Rep68 store concentration of the c	p
AAV-	COCCUTOCCATCTOCTCAACGTOCACCTOCCATCACTGTTTTTCACCCAATAAATGACTTAAACCACGTATCCCTGCCCATGGTTATCTTCCACATTACCTCACAATAAATGACTTCAACCACGTTATCTTCCACATTACCTCCACATTCACCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCTCACATTCACCTCTCTCTCTCACATTCACCTCTCTCTCTCTCTCACATTCACCTCTCTCTCTCTCTCTCACATTCACCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCACATTCACCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTCTC	2473
AAV-		2252
YYA-		
AAV-1	GOCATTCCCCLOTOGTCCGCACCTCCLACCCCCALACCCCCLALACCCLALACCCLALACCLACACCCCCGGGGTCCTTCCTGGCTACACGTACCTCGCACCCTTCLAC	2373
AAV- 3	.A.A.AA.ACA.OC.CCC.A.ACCA.AGCGCAGGGC.TAAAT	2373
***	,	2176
	•	
AAV.1	COLCTOCACA ACCOCCACCCCOTTCA ACCOCCACCACCACCACCACCACCACCACCACCACCACCA	2513
		3433
AAV-6		2478
	COCCADITICADE ACCUTE TO CALCALIA AND TETTT TO COCCAL CENTER CONCERCIANT CONCERCIANA ACCUSO THE TO CALCALIA ACCUTE TO CONCERCIANA ACCUSO TA LA CONCERCIANA ACCUSO TO CONCERCIANA	2632
<b>774-</b> 7		2613
AAV - 2		3618
	VP2	
AAV-1	VES - TOUR CONTROL OF THE PROPERTY OF THE PROP	2753
LAT-3	6AGG	2733
<b>117-6</b>	TGAC.T	****
	· VP3	
.AV.1	THE PROPERTY OF THE PROPERTY O	2173
		2832
M4-6		2050
	CCCCACCUACYCOCTAATOCCTCACGAAATTCCCACATTCCACATTCCACATCCCCCCACATCACCAC	2993
	m = = = = = = = = = = = = = = = = = = =	2973
MY-6		2978
MA-I	AACAANTCTCHATCCACTACAACAACAACAACAACAACAACAACAACAACA	3020
74-3	······································	3070
WA-0	••••••	
AV-1	CASSISTEMENTATIONS ATTOCOCCOMMENCATION ACTITICAL CITICAL CITICAL CITICAL CITICAL CONTRACT CANCELLA CON	3233
TA-3		3230
WA-R		****
AV-1	ACCIDENCEST TEAMORT TO TOTO ELECTROMISTACIA COTTO CONTROL CONT	1311
		3334
AV-4		3338
	TACCHIACOCTCUICALTOCOLOCOLOCCOTCOGOCOTTCATCCTTTTACTOCCTOGUATATTTCCCTTCTCACATOCTCACAACTATACCTTCACCTACACCTTTTAGA	1473
AV-1	Ç£Q	3450
AT-4		3458
AV-4	•	
FA-7	CLACTROCT TTCCACCACCTACOCCCCACACCCCCCACCCCCCATGLATGLATCCTCTCATCACCCATATACCCTCTATTACCTCTAACACACACACACACACACACACACACACACACACACACA	3373
FA-3	.Ct	96/T 1672
7A-4	··G···································	
AV. 1	ALCIAGIACTICITOTITIACCOMOGNICICCIACTOCCATGICTOTICACCCCAAAACTOCCTACCTOCACCCGCTACCCCCCCCCC	713
	G GEORGE B 1 WEST AND COOKER GARD. A TOOG. A.C T. T. CO.C	
AV-6		698
	•	
7A-J	ANCICCIATITITACCIOGRACTOGRACTICARARIATARACCICATOGRACTICATCATCATCATCATCATCATCATOCACCICATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATATOGRACTICATAT	014
<b>LA-3</b>	TG.A.ACT.GA.CG.CCC	018

## FIG IC

	AGCOGGGCATGATTITTOGGAAAAGAGACCCCCGGAGCTTCAAACACTOCATTOGACGATGATTACAGACGAAGAGAGAAAAATTAAAGCCACTAGACCCTGGAAGATTT 3:G.TC.C.C.G.GC.JG.T.A.AGAAAA.TGCGACAT.A.G	
	OGGACGGTGGCAGTCAATTTCCAGACCAGCAGCACAACAACCCTGCGACGGGAATGTGCATGCTATGGAGAGATAACTGGAGTGGGAACAAAGACAACAACAA.C.TTC.T.AT.TAC.CCCA.G.C.G.G.G.G	
	ATTTOGGCCAAAATTCCTCACACAGATGGACACTTTCACCCGTCTCCTC.TTATGGCCGGCTTTGGACTCAAGAACCCGCCTCCTCACATCCTCATCAAAAACACGCCTGTTCCTGCCAAA 41CAGAGCGAGCGAGCGAGCGAGGG	
	CCTCCCCCCCCCCAGTTTTCACCTACAAAGTTTCCTTCATCACCCCAATACTCCACCCCCA-AGTGLGTGTCCCAAAATTCAATGCCACCTCCCAGAACCACCTCCAATCC 41T.A.CACC. CAGT00	
	COLAGOCAGIACACTICALTIATOCALAGICTOCCAA-COTTGATTTACTGTUGACAACAATOCACTTATACCCACCCCCATTGCCACCCCGTTACCCCCCCTTTACCCCCCCC	
AV-1	VP1-3 stop       PolyA signal         ANTICOTOTTANICATEMACCOSTRUCTURE CONTROLLEGISTIC CONTRO	
	TTCUCCATAAAACACTTACGTCATCCCCCC LAGCCCCLBytynafttgcccadlccctctctgcgcgctcgctcggtgggggcgggcagugagsgctctgccgctcggtggggccagugagsgctctgccgctcggtggggccggcagugagsgctctgccgctcggtggggccggcagugagsgctctgccgctcggtgggggccggcagugagsgctctgccggtggggactgctcggctcggtcggtcggtcg	
AV-2	Egyttcqcagyctcaccgagcgagcgagcgcagagagggagtgyycaa 4718co.g.gct.gt	

	α,
	E
	<del>,  </del>
	7
•	A
	A








FIG. 3A










Group	1	2	3	4	5	6
Vector1 - a1AT						
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2



Group	1	2	3	4	5	6
Vector1 - α1AT	AAV2	AAV1	PBS	PBS	AAV2	AAV1
Vector2-EPO	AAV2	AAV1	AAV2	AAV1	AAV1	AAV2

## SEQUENCE LISTING

<110> Wilson, James M. Xiao, Weidong The Trustees of the University of Pennsylvania

<120> Adeno-Associated Virus Serotype I Nucleic Acid Sequences, Vectors and Host Cells Containing Same

<130> GNVPN.031PCT

<140>

<141>

<150> 60/107,114

<151> 1998-11-05

<160> 20

<170> PatentIn Ver. 2.0

<210> 1

<211> 4718

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (335)..(2206)

<220>

<221> CDS

<222> (2223).. (4430)

<400> 1

ttgcccactc cctctctgcg cgctcgctcg ctcggtgggg cctgcggacc aaaggtccgc 60 agacggcaga getetgetet geeggeeeca eegagegage gagegegeag agagggagtg 120 ggcaactcca tcactagggg taatcgcgaa gcgcctccca cgctgccgcg tcagcgctga 180 cgtaaattac gtcatagggg agtggtcctg tattagctgt cacgtgagtg cttttgcgac 240 attttgcgac accacgtggc catttagggt atatatggcc gagtgagcga gcaggatctc 300 cattttgacc gcgaaatttg aacgagcagc agcc atg ccg ggc ttc tac gag atc 355

Met Pro Gly Phe Tyr Glu Ile

gtg Val	atc Ile	aag Lys 10	gtg Val	ccg Pro	agc Ser	gac Asp	ctg Leu 15	gac Asp	gag Glu	cac His	ctg Leu	ccg Pro 20	ggc Gl <b>y</b>	att Ile	tct. Ser	403
gac Asp	tcg Ser 25	ttt Phe	gtg Val	agc Ser	tgg Trp	gtg Val 30	gcc Ala	gag Glu	aag Lys	gaa Glu	tgg Trp 35	gag Glu	ctg Leu	ccc Pro	ccg Pro	451
gat Asp 40	tct Ser	gac Asp	atg Met	gat Asp	ctg Leu 45	aat Asn	ctg Leu	att Ile	<b>g</b> ag Glu	cag Gln 50	gca Ala	ccc Pro	ctg Leu	acc Thr	gtg Val 55	499
gcc Ala	gag Glu	aag Lys	ctg Leu	cag Gln 60	cgc Arg	gac Asp	ttc Phe	çtg Leu	gtc Val 65	caa Gln	tgg Trp	cgc Arg	cgc Arg	gtg Val 70	agt Ser	547
aag Lys	gcc Ala	ccg Pro	gag Glu 75	gcc Ala	ctc Leu	ttc Phe	ttt Phe	gtt Val 80	cag Gln	ttc Phe	gag Glu	aag Lys	ggc Gly 85	gag Glu	tcc Ser	595
													aaa Lys			643
													gtg Val			691
atc Ile 120	tac Tyr	cgc Arg	Gly gg <b>g</b>	atc Ile	gag Glu 125	ccg Pro	acc Thr	ctg Leu	ccc Pro	aac Asn 130	tgg Trp	ttc Phe	gcg Ala	gtg Val	acc Thr 135	739
aag Lys	acg Thr	cgt Arg	aat Asn	ggc Gly 140	gcc Ala	gga Gly	ggg ggg	el y ggg	aac Asn 145	aag Lys	gtg Val	gtg Val	gac Asp	gag Glu 150	tgc Cys	787
tac Tyr	atc Ile	ccc Pro	aac Asn 155	tac Tyr	ctc Leu	ctg <b>Leu</b>	ccc Pro	aag Lys 160	act Thr	cag Gln	ccc Pro	gag Glu	ctg Leu 165	cag Gln	tgg Trp	835
gcg Ala	tgg Trp	act Thr 170	aac Asn	atg Met	gag Glu	gag Glu	tat Tyr 175	ata Ile	agc Ser	gcc Ala	tgt Cys	ttg Leu 180	aac Asn	ctg Leu	gcc Ala	883
gag Glu	cgc Arg 185	aaa Lys	cgg Arg	ctc Leu	gtg Val	gcg Ala 190	cag Gln	cac	ctg Leu	acc Thr	cac His 195	gtc Val	agc Ser	cag Gln	acc Thr	931

-	Glu			: aag Lys		Asn	•					•			_	979
				acc Thr 220					_		-	-	,,,	,,,	_	1027
	-			atc				_						-	_	1075
-	-			tcc Ser			-	- T			-			•		1123
-	-	-	_	gac Asp				_		-		•				1171
		_		ctg Leu	-			_	_		•	-				1219
	-			cgc Arg 300		-		-				-		-		1267
_			_	ttt Phe				-	_					_	-	1315
				ctg Leu			-	-				-				1363
	-	-		gcc Ala		_						-	-			1411
				ttt Phe				-	-	-	_	-	-	-		1459
				ggc Gly 380	_	_	-	Ala	_	_			Ser	-	-	1507

-				y Gly	-		-	-	g Va				-	s Ly	g tcg s Ser	1555
			n Ile					o Val					: Ası		c aac r Asn	1603
		a Ala					Ası	-				Glu		-	g cag n Gln	1651
						Phe					Thr	-	_	-	gag Glu 455	1699
His	Asp	Phe	: Gly	aag Lys 460	Val	Thr	Lys	Gln	Glu 465	Val	Lys	Glu	Phe	Phe 470	Arg	1747
Trp	Ala	Gln	Asp 475	cac His	Val	Thr	Glu	Val 480	Ala	His	Glu	Phe	Tyr 485	Val	Arg	1795
Lys	Gly	Gly 490	Ala	aac Asn	Lys	Arg	Pro 495	Ala	Pro	Asp	Asp	Ala 500	Asp	Lys	Ser	1843
Glu	Pro 505	Lys	Arg	gcc Ala	Cys	Pro 510	Ser	Val	Ala	Asp	Pro 515	Ser	Thr	Ser	Asp	1891
Ala 520	Glu	Gly	Ala	ccg Pro	Val 525	Asp	Phe	Ala	Asp	Arg 530	Tyr	Gln	Asn	Lys	Cys 535	1939
Ser	Arg	His	Ala	ggc Gly 540	Met	Leu	Gln	Met	Leu 545	Phe	Pro	Cys	Lys	Thr 550	Cys	1987
Glu .	Arg	Met	Asn 555	cag Gln	Asn	Phe	Asn	Ile 560	Cys	Phe	Thr	His	Gly 565	Thr	Arg	2035
	Cys			tgc Cys		Pro					Ser					2083

-	_	Arg		tat Tyr			Leu									2131
	-			att Ile		Cys					Leu					2179
_	-	-	_	gtt Val 620				taa	atg	actt	aaa	ccag	M		ct gcc la Ala	2231
-				cca Pro	_											2279
-	_			gac Asp												2327
9		_	_	gac Asp					_						-	2375
				ttc Phe 680				-			-					2423
	-	-		gcc Ala												2471
			-	aat Asn												2519
				ctg Leu												2567
_	-	-		cag Gln	_	-	_		_							2615
_		-		gct Ala 760	-	-		Pro		•		-	-	-		2663

				Glu			Ser				ggc Gly	<b>271</b> 1
_	_		Ala			Asn			act Thr 800			2759
		Val		_					cca Pro			2807
-	-								ggc Gly			2855
-	_			_					gcc Ala			2903
									atc Ile			2951
	-			_					ctc Leu 880			2999
									cac His			3047
									ttc Phe			3095
									aat Asn			3143
									caa Gln			3191
					Val			Asn	aac Asn 960			3239

acq qtt caa	a gtc ttc tcg	gac tog gad	tac cag ct	t.ccg tac gtc	ctc 3287
	-			u Pro Tyr Val	
965		970	97:		
303		3.0			
		tac etc cct	eca tte ce	g gcg gac gtg	ttc 3335
				o Ala Asp Val	
-	_			Ala Ash var	995
980	985		990		333
					2202
_				ggc agc caa	
Met Ile Pro		Tyr Leu Thr		Gly Ser Gln	Ala
	1000		1005	1010	
				cct tct cag	
Val Gly Arg	Ser Ser Phe	Tyr Cys Leu	Glu Tyr Phe	Pro Ser Gln	Met
	1015	1020		1025	
			•		
ctg aga acg	ggc aac aac	ttt acc ttc	agc tac acc	ttt gag gaa	gtg 3479
Leu Arg Thr	Gly Asn Asn	Phe Thr Phe	Ser Tyr Thr	Phe Glu Glu	Val
1030		1035		1040	
cct ttc cac	age age tac	gcg cac agc	cag agc ctg	gac cgg ctg	atg 3527
				Asp Arg Leu	
1045	_	1050	1055		
1045	•	2000	200		
aat oot oto	atc dac caa	tac ctg tat	tac ctg aac	aga act caa	aat 3575
				Arg Thr Gln	
	1065	Tyl Deu Tyl	1070		1075
1069	1005		1070	•	1075
	ost see 633	226 226 626	***	. agc cgt ggg	tct 3623
_					
Gin Ser Giy				Ser Arg Gly	261
	1080		1085	1090	
					tat 3671
	-	_		cct gga ccc	
_			Asn Trp Leu	Pro Gly Pro	Cys
	1095	1100		1105	
,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	2 2 2		-	aac aac aac	
Tyr Arg Gln	Gln Arg Val	Ser Lys Thr	Lys Thr Asp	Asn Asn Asn	Ser
1110		1115		1120	
aat ttt acc	tgg act ggt	gct tca aaa	tat aac ctc	aat ggg cgt	gaa 3767
Asn Phe Thr	Trp Thr Gly	Ala Ser Lys	Tyr Asn Leu	Asn Gly Arg	Glu
1125	1	130	1135		
tcc atc atc	aac cct ggc	act gct atg	gcc tca cac	aaa gac gac	gaa 3815
			_	Lys Asp Asp	
1140	1145		1150		.155
_ <del></del> -					

															agc Ser	3863
	,			1160			1		1165			-		1170		
-															gaa	3911
Ala	Gly		Ser	Asn	Thr	Ala			Asn	Val	Met			Asp	Glu	
			1175					1180					1185			
	_		aaa	-												<b>39</b> 59
Glu			Lys	Ala	Thr			Val	Ala	Thr			Phe	Gly	Thr	
		1190					1195				•	1200				
			aat													4007
Val	Ala	Val	Asn	Phe	Gln	Ser	Ser	Ser	Thr			Ala	Thr	Gly	Asp	
1	1205					1210				-	1215					
gtg	cat	gct	atg	gga	gca	tta	cct	ggc	atg	gtg	tgg	caa	gat	aga	gac	4055
Val	His	Ala	Met	Gly	Ala	Leu	Pro	Gly	Met	Val	Trp	Gln	Asp			
1220	)			]	1225				1	L230					1235	
gtg	tac	ctg	cag	ggt	ccc	att	tgg	gcc	aaa	att	cct	cac	aca	gat	gga	4103
Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp	Ala	Lys	Ile	Pro	His	Thr	Asp	Gly	
			1	1240					1245				-	1250		
cac	ttt	cac	ccg	tct	cct	ctt	atg	ggc	ggc	ttt	gga	ctc	aag	aac	ccg	4151
His	Phe	His	Pro	ser	Pro	Leu	Met	Gly	Gly	Phe	Gly			Asn	Pro	
		]	1255				1	L260				1	1265			
			atc													4199
Pro	Pro	Gln	Ile	Leu	Ile	Lys	Asn	Thr	Pro	Val			Asn	Pro	Pro	
	1	1270				1	1275				1	.280				
			tca	_		-										4247
Ala	Glu	Phe	Ser	Ala	Thr	Lys	Phe	Ala	Ser			Thr	Gln	Tyr	Ser	
1	.285				1	.290				1	1295					
aca	gga	caa	gtg	agt	gtg	gaa	att	gaa	tgg	gag	ctg	cag	aaa	gaa	aac	4295
Thr	Gly	Gln	Val	Ser	Val	Glu	Ile	Glu	Trp	Glu	Leu	Gln	Lys			
1300	)			1	.305				1	.310				•	1315	
agc	aag	cgc	tgg	aat	ccc	gaa	gtg	cag	tac	aca	tcc	aat	tat	gca	aaa	4343
Ser	Lys	Arg	Trp	Asn	Pro	Glu	Val	Gln	Tyr	Thr	ser	Asn	Tyr	Ala	Lys	
			1	320				1	325				1	L <b>3</b> 30		
tct	gcc	aac	gtt	gat	ttt	act	gtg	gac	aac	aat	gga	ctt	tat	act	gag	4391
Ser.	Ala	Asn	Val	Asp	Phe	Thr	Val	Asp	Asn	Asn	Gly	Leu	Tyr	Thr	Glu	
		1	.335				. 1	340				1	345			

cct cgc ccc att ggc acc cgt tac ctt acc cgt ccc ctg taattacgtg. 4440
Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
1350 1355 1360

ttaatcaata aaccggttga ttcgtttcag ttgaactttg gtctcctgtc cttcttatct 4500
tatcggttac catggttata gcttacacat taactgcttg gttgcgcttc gcgataaaag 4560
acttacgtca tcgggttacc cctagtgatg gagttgccca ctccctctct gcgcgctcgc 4620
tcgctcggtg gggcctgcgg accaaaggtc cgcagacggc agagctctgc tctgccggcc 4680
ccaccgagcg agcgagcgcg cagagaggga gtgggcaa 4718

<210> 2

<211> 623

<212> PRT

<213> AAV-1

<400> 2

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp
1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 

Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415

Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430

Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe
435
440
445

Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460

Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 535 540

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 615 620

<210> 3

<211> 736

<212> PRT

<213> AAV-1

_	$\sim$	0>	- 3

Met	Ala	Ala	Asp	Gly	Tyr	Leu	Pro	Asp	Trp	Leu	Glu	Asp	Asn	Leu	Ser
1				5					10					15	

- Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30
- Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45
- Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro 50 55 60
- Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80
- Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95
- Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
  100 105 110
- Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125
- Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140
- Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly 145 150 155 160
- Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175
- Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190
- Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly 195 200 205
- Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220
- Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240
- Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu

245 250 255

Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270

Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285

His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300

Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320

Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335

Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350

Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365

Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380

Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400

Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe 405 410 415

Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430

Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg 435 440 445

Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 455 460

Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480

Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495

Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn

500 505 510

Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525

Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540

Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560

Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg 565 570 575

Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590

Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln 595 600 605

Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 615 620

Thr Asp Gly His Phe His Pro Ser Pro Leu Mot Gly Gly Phe Gly Leu 625 635 640

Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655

Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670

Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685

Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 4

<211> 1872

<212> DNA

<213> AAV-1

<220>
<221> CDS
<222> (1)..(1869)

<400> 4

atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp

1 5 10 15

gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag 96
Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu
20 25 30

aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att 144
Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile
35 40 45

gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

gtc caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt 240
Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val
65 70 75 80

cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag 288
Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu
85 90 95

acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att 336
Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile
100 105 110

agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu

115

120

125

ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag 480 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

act cag ccc gag ctg cag tgg gcg tgg act aac atg gag gag tat ata 528

	•																
	Thr	Gln	Pro	Glu	Leu 165		Тгр	Ala	Trp	Thr 170		Met	Glu	Glu	Туr 175	Ile	* *
	-	_			aac Asn												576
	_			Val	agc Ser	_							_			aac , Asn	624
				-	gcg Ala												672
					GJ y ggg												720
	-			-	gag Glu 245	-	_	_	_						_	-	768
			-		tcc Ser	-		-	-	-	_	-		-		_	816
		_	-	_	acc Thr					_		-	-			-	864
	-		-	-	att Ile				-			_		_		-	912
				_	cct Pro	-		-			-					_	960
	-				ggg Gly 325	_	-		Thr			_			_	_	1008
		_		_	acc Thr			Ala	_	_		_		-			1056
1	ttc	tac	ggc	tgc	gtc	aac	tgg	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	1104

Phe	Tyr	Gly 355		Val	Asn	Trp	Thr 360		Glu	Asn	Phe	9ro 365		Asn	Asp	
_	-	Asp		atg Met			Trp					Lys				1152
_	_			tcc Ser							Gly					1200
	-			tgc Cys 405												1248
	_			aac Asn												1296
				cac His												1344
_				cgt Arg												1392
_	-			ttc Phe												1440
				tac Tyr 485												1488
-	_	_	, ,	gat Asp		_			_		-	-			-	1536
gcg Ala	-		_	acg Thr		_		_		-	_		-		_	1584
gac Asp					Lys	-		-				Met		-	-	1632
ctg	ttt	ccc	tgc	aag	aca	tgc	gag	aga	atg	aat	cag	aat	ttc	aac	att	1680

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 5.55 550 545 tgc ttc acg cac ggg acg aga gac tgt tca gag tgc ttc ccc ggc gtg 1728 Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 570 565 tea gam tet cam deg gte gte aga mag agg meg tat egg mam ete tgt 1776 Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 585 580 gcc att cat cat ctg ctg ggg cgg gct ccc gag att gct tgc tcg gcc 1824 Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 600 . 595 tgc gat ctg gtc aac gtg gac ctg gat gac tgt gtt tct gag caa taa 1872 Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 615 <210> 5 <211> 623 <212> PRT <213> AAV-1 <400> 5 Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 15 5. Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 55 Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 80 . 70 75 Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 105 110 100 Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu

115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys
145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190

Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 235

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255

Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270

Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285

Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300

Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315 320

Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335

Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350

Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365

Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala

370 375 380

Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400

- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445
- Glu Leu Thr Arg Arg Leu Glu His Asp Phe Gly Lys Val Thr Lys Gln 450 455 460
- Glu Val Lys Glu Phe Phe Arg Trp Ala Gln Asp His Val Thr Glu Val 465 470 475 480
- Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala 485 490 495
- Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val 500 505 510
- Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 515 520 525
- Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 530 535 540
- Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 545 550 555 560
- Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 565 570 575
- Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 580 585 590
- Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 595 600 605
- Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 610 620

<210> 6 <211> 1641 <212> DNA <213> AAV-1 <220> <221> CDS <222> (1)..(1638) <400> 6 atg ccg ggc ttc tac gag atc gtg atc aag gtg ccg agc gac ctg gac Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp 5 gag cac ctg ccg ggc att tct gac tcg ttt gtg agc tgg gtg gcc gag Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 aag gaa tgg gag ctg ccc ccg gat tct gac atg gat ctg aat ctg att Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 gag cag gca ccc ctg acc gtg gcc gag aag ctg cag cgc gac ttc ctg 192 Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu gto caa tgg cgc cgc gtg agt aag gcc ccg gag gcc ctc ttc ttt gtt Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 70 75 288 cag ttc gag aag ggc gag tcc tac ttc cac ctc cat att ctg gtg gag Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 336 acc acg ggg gtc aaa tcc atg gtg ctg ggc cgc ttc ctg agt cag att Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 agg gac aag ctg gtg cag acc atc tac cgc ggg atc gag ccg acc ctg 384 Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125 ccc aac tgg ttc gcg gtg acc aag acg cgt aat ggc gcc gga ggg ggg 432 Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly 130 135 aac aag gtg gtg gac gag tgc tac atc ccc aac tac ctc ctg ccc aag 480 Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys

acc acg ggc aag acc aac atc gcg gaa gcc atc gcc cac gcc gtg ccc Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro

340 345 350

tto	: tac	ggc	: tgc	gto	aac	tgg	acc	: aat	gag	aac	: ttt	ccc	: tto	aat	t gat	1104	
Phe	Tvr	Glv	CVS	· Val	l Ası	Tre	Thr	Asr	Glu	Asn	Phe	Pro	Phe	e Ası	Asp		
	- 3	355	_				360					365					
		333	,				300	,				300	,				
tgc	gtc	gac	aag	ato	gto	, atc	: tgg	tgg	gag	gag	ggc	aag	ato	acq	gcc	1152	
Cys	Val	Asp	Lys	Met	: Val	Ile	Trp	Trp	Glu	Glu	Gly	Lys	Met	Thi	Ala		
-	370	_				375					380	) =					
											•						
-	-				-	_	-					-	-		, cgc	1200	
Lys	Val	Val	Glu	Ser	Ala	Lys	Ala	Ile	Leu	Gly	Gly	Ser	Lys	Val	Arg		
385					390	•				<b>39</b> 5					400		
ata	gac	caa	aaσ	tac	aad	tca	tee	acc.	cad	atc	gac	ccc	acc	CCC	gtg	1248	
	-		_	_	_	_		-	-		-					1210	
Val	Asp	GIII	гЛЭ	-	-	ser	ser	AIA		116	Asp	PIO	Inr		Val		
				405					410					415			
atc	gtc	acc	tcc	aac	acc	aac	atg	tgc	gcc	gtg	att	gac	qqq	aac	agc	1296	
Tle	Val	Thr	Ser	Asn	Thr	Asn	Met	Cvs	Ala	Val	Tle	Asp	Glv	Asn	Ser		
			420					425					430		502		
			420					423					430				
acc	acc	ttc	gag	cac	cag	cag	ccg	ttg	cag	gac	cgg	atg	ttc	aaa	ttt	1344	
Thir	Thr	Phe	Glu	His	Gln	Gln	Pro	Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe		
		435					440					445					
ma a	ctc	200	cac	cat	cta	a a a	cat	asc.	+++	aac	2 2 7	ata	202	224	cag	1392	
														_	_	1392	
GIU		Thr	Arg	Arg	Leu	Glu	HIS	Asp	Pne	GIA	-	vaı	Thr	Lys	GIn		
	450					455					460						
gaa	gtc	aaa	gag	ttc	ttc	cgc	tyg	gcg	cag	gat	cac	gtg	acc	gag	gtg	1440	
Glu	Val	Lys	Gìu	Phe	Phe	Arg	Trp	Ala	Gln	Asp	His	Val	Thr	Glu	Val		
465		•			470	-	•			475					480		
										1,5					400		
						aga									-	1488	
Ala	His	Glu	Phe	Tyr	Val	Arg	Lys	Gly	Gly	Ala	Asn	Lys	Arg	Pro	Ala		
				485					490					495			
ccc	nat	aac	aca	αat	aaa	agc	a a a	ccc	224	caa	acc	tac	CCC	+ = =	ata	1536	
									_			-			_	1550	
PIO	ASP	ASp		Asp	ràz	Ser	GIU		Lys	Arg	Ата	cys		ser	vaı		
			500					505					510				
gcg	gat	cca	tcg	acg	tca	gac	gcg	gaa	gga	gct	ccq	gtq	gac	ttt	gcc	1584	
			_	_		Asp		_		-	_		-		-		
	_	515				-	520		1				- ~P		. <del></del> u		
		J I J					J20					525					
gac	agg	tat	ggc	tgc	cga	tgg	tta	tct	tcc	aga	ttg	gct	cga	gga	caa	1632	
Asp	Arg	Tyr	Gly	Cys	Arg	Trp	Leu	Ser	Ser .	Arg	Leu	Ala	Arg	Gly	Gln		

530 535 540

cct ctc tga 1641 .

Pro Leu 545

<210> 7

<211> 546

<212> PRT

<213> AAV-1

<400> 7

Met Pro Gly Phe Tyr Glu Ile Val Ile Lys Val Pro Ser Asp Leu Asp
1 5 10 15

Glu His Leu Pro Gly Ile Ser Asp Ser Phe Val Ser Trp Val Ala Glu 20 25 30

Lys Glu Trp Glu Leu Pro Pro Asp Ser Asp Met Asp Leu Asn Leu Ile 35 40 45

Glu Gln Ala Pro Leu Thr Val Ala Glu Lys Leu Gln Arg Asp Phe Leu 50 55 60

Val Gln Trp Arg Arg Val Ser Lys Ala Pro Glu Ala Leu Phe Phe Val 65 70 75 80

Gln Phe Glu Lys Gly Glu Ser Tyr Phe His Leu His Ile Leu Val Glu 85 90 95

Thr Thr Gly Val Lys Ser Met Val Leu Gly Arg Phe Leu Ser Gln Ile 100 105 110

Arg Asp Lys Leu Val Gln Thr Ile Tyr Arg Gly Ile Glu Pro Thr Leu 115 120 125

Pro Asn Trp Phe Ala Val Thr Lys Thr Arg Asn Gly Ala Gly Gly Gly 130 135 140

Asn Lys Val Val Asp Glu Cys Tyr Ile Pro Asn Tyr Leu Leu Pro Lys 145 150 155 160

Thr Gln Pro Glu Leu Gln Trp Ala Trp Thr Asn Met Glu Glu Tyr Ile 165 170 175

Ser Ala Cys Leu Asn Leu Ala Glu Arg Lys Arg Leu Val Ala Gln His 180 185 190 Leu Thr His Val Ser Gln Thr Gln Glu Gln Asn Lys Glu Asn Leu Asn 195 200 205

- Pro Asn Ser Asp Ala Pro Val Ile Arg Ser Lys Thr Ser Ala Arg Tyr 210 215 220
- Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 225 230 235 240
- Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 245 250 255
- Ser Asn Ser Arg Ser Gln Ile Lys Ala Ala Leu Asp Asn Ala Gly Lys 260 265 270
- Ile Met Ala Leu Thr Lys Ser Ala Pro Asp Tyr Leu Val Gly Pro Ala 275 280 285
- Pro Pro Ala Asp Ile Lys Thr Asn Arg Ile Tyr Arg Ile Leu Glu Leu 290 295 300
- Asn Gly Tyr Glu Pro Ala Tyr Ala Gly Ser Val Phe Leu Gly Trp Ala 305 310 315 320
- Gln Lys Arg Phe Gly Lys Arg Asn Thr Ile Trp Leu Phe Gly Pro Ala 325 330 335
- Thr Thr Gly Lys Thr Asn Ile Ala Glu Ala Ile Ala His Ala Val Pro 340 345 350
- Phe Tyr Gly Cys Val Asn Trp Thr Asn Glu Asn Phe Pro Phe Asn Asp 355 360 365
- Cys Val Asp Lys Met Val Ile Trp Trp Glu Glu Gly Lys Met Thr Ala 370 375 380
- Lys Val Val Glu Ser Ala Lys Ala Ile Leu Gly Gly Ser Lys Val Arg 385 390 395 400
- Val Asp Gln Lys Cys Lys Ser Ser Ala Gln Ile Asp Pro Thr Pro Val 405 410 415
- Ile Val Thr Ser Asn Thr Asn Met Cys Ala Val Ile Asp Gly Asn Ser 420 425 430
- Thr Thr Phe Glu His Gln Gln Pro Leu Gln Asp Arg Met Phe Lys Phe 435 440 445

	450					455					460		Thr			
GLu 465	Val	Lys	Glu	Phe	Phe 470	Arg	Trp	Ala	Gln	Asp 475	His	Val	Thr	Glu	Val 480	
Ala	His	Glu	Phe	Tyr 485	Vāl	Arg	Lys	Gly	Gly 490	Ala	Asn	Lys	Arg	Pro 495	Ala	
Pro	Asp	Asp	Ala 500	Asp	Lys	Ser	Glu	Pro 505	Lys	Arg	Ala	Cys	Pro 510	Ser	Val	
		515					520					525	Asp			
Asp	Arg 530	Tyr	Gly	Cys	Arg	Trp 535	Leu	Ser	Ser	Arg	Leu 540	Ala	Arg	Gly	Gln	
Pro	Leu															
545					•											
<210	)> 8															
<211	L> 12	200														
-	2> D1															
	3> A7	\V-1												•		
<220	)> L> CI	ne														
		L) (	(1197	7)												
<400																
atg	gag	ctg	gtc	ggg	tgg	ctg	gtg	gac	cgg	ggc	atc	acc	tcc	gag	aag -	48
Met	Glu	Leu	Val		Trp	Leu	Val	Asp		Gly	Ile	Thr	Ser		Lys	
1				5					10					15		
cag	tgg	atc	cag	gag	gac	cag	gcc	tcg	tac	atc	tcc	ttc	aac	gcc	gct	96
													Asn			
			20					25					30			
tcc	aac	tcq	cgg	tcc	cag	atc	aag	gcc	gct	ctg	gac	aat	gcc	ggc	aag	144
													Al.a			
		35					40					45				
atc	atσ	aca	cta	acc	aaa	tcc	aca	ccc	gac	tac	ctg	gta	ggc	ccc	gct	192
													Gly			
	50					55			-	_	60					

ccq	ccc	gcg	gac	att	aaa	acc	aac	cgc	atc	tac	cgc	atc	ctg	gag	ctg	240
Pro	Pro	Ala	Asp	Ile	Lvs	Thr	Asn	Arg	Ile	Tyr	Arg	Ile	Leu	Glu	Leu	
		•			70			_		75					80	
65					,,											
										~+~		ctc	aac	taa	acc	288
aac	ggc	tac	gaa	cct	gcc	tac	gcc	ggc	- CCC	gtc	-1	-	ggc	cyy	71-	200
Asn	Gly	Tyr	Glu	Pro	Ala	Tyr	Ala	Gly	Ser	Val	Phe	Leu	Gly		Ala	
				85					90					95		
			++-	~~~	220	cac	220	acc	atc	taa	cta	ttt	ggg	cca	acc	336
cag	aaa	agg	-1	999	aay	na-	Dan	mb=	Tlo	423	Tan	Dhe	Glu	Pro	Δla	
Gln	Lys	Arg	Pne	GTÀ	ràz	Arg	Asn		116	rrb	Бец	LIIC	Gly	110	,,,,,	
			100					105					110			
					•											
acc	acq	aac	aaq	acc	aac	atc	qcq	gaa	gcc	atc	gcc	cac	gcc	gtg	CCC	384
mb -	mbs	23.	Tuc	Thr	Nen	Tle	Δla	Glu	Ala	Tle	Ala	His	Ala	Val	Pro	
Thr	THE		гуз	1111	Mali	110						125				
		115					120					125				
ttc	tac	ggc	tgc	gtc	aac	tgg	acc	aat	gag	aac	ttt	ccc	ttc	aat	gat	432
Phe	Tvr	Gly	Cys	Val	Asn	Trp	Thr	Asn	Glu	Asn	Phe	Pro	Phe	Asn	Asp	
	130	•	•			135					140					
	130															
						25.5	+ ~ ~	taa	n a n	asa	aac	ааσ	atg	acd	gcc	480
tgc	gtc	gac	aag	atg	gra	acc	- cgg	- cgg	gay	gag	99°	Tue	Wat	Th-	מות	
Cys	Val	Λsp	Lys	Met	Val	Ile	Trp	Trp	GLu		GIA	гAг	Met	THE		
145					150					155					160	
aan	atc	ata	σασ	tcc	qcc	aaq	qcc	att	ctc	ggc	ggc	agc	aag	gtg	cgc	528
1	Val	V-1	Glu	Ser	Δla	Lvs	Ala	Ile	Leu	Glv	Glv	Ser	Lys	Val	Arg	
гÀз	Vai	Val	GIU		AIU	Dy D			170	,	,		-	175	_	
				165					170					1.0		
																520
gtg	gac	caa	aag	tgc	aag	tcg	tcc	gcc	cag	atc	gac	ccc	acc	CCC	gtg	576
Val	Asp	Gln	Lys	Cys	Lys	Ser	Ser	Ala	Gln	Ile	Asp	Pro	Thr	Pro	Val	
	-		180					185					190			
	4 -		•		200	226	250	tac	acc	ata	att	gac	ggg	aac	aσc	624
atc	gtc	acc	tee	aac	acc	aac	ary	cyc	31-	909	71-	7	233	) co	Ser	
Ile	Val	Thr	Ser	Asn	Thr	Asn		Cys	ALA	var	116		Gly	ASII	SCI	
		195					200					205				
acc	acc	ttc	gag	cac	cag	cag	ccg	ttg	cag	gac	cgg	atg	ttc	aaa	ttt	672
The	Thr	Dhe	Glu	His	Gln	Gln	Pro	Leu	Gln	Asp	Arg	Met	Phe	Lys	Phe	
1111		1110				215				•	220					
	210					210										
												_+-			<b>~~~</b>	720
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt	ggc	aag	gcg	aca	aag	cag	120
Glu	Leu	Thr	Arg	Arg	Leu	Glu	His	Asp	Phe	Gly	Lys	Val	Thr	Lys		
225					230					235					240	
	- t-		~~~	++~	++~	cac	taa	aca	cad	gat	cac	ata	acc	σασ	ata	768
gaa	gcc	add	yay	21.	חויי	n	m	.y~y	C1 =	Der.	uic	Val	The	G111	Val	
Glu	Val	Lys	GIU		rne	Arg	rrp	MTg		νsδ	urs	val	Thr			
				245					250					255		

-		gag Glu		Tyr	-	-			Gly	-				816
	-	gac Asp 275		-				Pro						864
	-	cca Pro	_	_		_		_		-	_	 _	_	912
-		tac Tyr												960
,		ccc Pro	-	-		-	-	•	-		_			1008
-		acg Thr			_	-	-	-			-		 	1056
	•	tct Ser 355		-	-	-	-	-		-			_	1104
-		cat His		_		-								1152
-	-	ctg Leu	-		-	-	_	-	-	-	-		taa	1200

<210> 9

<211> 399

<212> PRT

<213> AAV-1

<400> 9

Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys

1 5 10 15

Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala

30

			20					25	•				30		
Ser	Asn	Ser 35		Ser	Gln	Ile	Lys 40		Ala	Leu	Asp	Asn 45		Gly	Ly
Ile	Met 50	Ala	Leu	Thr	Lys	Ser 55		Pro	Asp	тyr	Leu 60		Gly	Pro	Ala
Pro 65	Pro	Ala	Asp	Ile	Lys 70	Thr	Asn	Arg	Ile	Tyr 75	Arg	Ile	Leu	Glu	Lei 8
Asn	Gly	Туr	Glu	Pro 85	Ala	туг	Ala	Gly	Ser 90		Phe	Leu	Gly	Trp 95	Ala
Gln	Lys	Arg	Phe 100	Gly	Lys	Arg	Asn	Thr 105	Ile	Trp	Leu	Phe	Gly 110	Pro	Ala
Thr	Thr	Gly 115	Lys	Thr	Asn	Ile	Ala 120	Glu	Ala	Ile	Ala	His 125	Ala	Val	Pro
	130					135					140			Asn	
145					150					155				Thr	160
Lys	Val	Val	Glu	Ser 165	Ala	Lys	Ala	Ile	Leu 170	Gly	Gly	Ser	Lys	Val 175	Arg
	•		180	-	-			185					190	Pro	
Ile	Val	Thr 195	Ser	Asn	Thr	Asn	Met 200	Cys	Ala	Val	Ile	Asp 205	Gly	Asn	Ser
Thr	Thr 210	Phe	Glu	His	Gln	Gln 215	Pro	Leu	Gln	Asp	Arg 220	Met	Phe	Lys	Phe
Glu 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys	Val	Thr	Lys	Gln 240
Glu	Val	Lys	Glu	Phe 245	Phe	Arg	Trp	Ala	Gln 250	Asp	His	Val	Thr	Glu 255	Val

260

Ala His Glu Phe Tyr Val Arg Lys Gly Gly Ala Asn Lys Arg Pro Ala

Pro Asp Asp Ala Asp Lys Ser Glu Pro Lys Arg Ala Cys Pro Ser Val

285

275

280

Ala Asp Pro Ser Thr Ser Asp Ala Glu Gly Ala Pro Val Asp Phe Ala 300 295 290

Asp Arg Tyr Gln Asn Lys Cys Ser Arg His Ala Gly Met Leu Gln Met 315 310

Leu Phe Pro Cys Lys Thr Cys Glu Arg Met Asn Gln Asn Phe Asn Ile 330

Cys Phe Thr His Gly Thr Arg Asp Cys Ser Glu Cys Phe Pro Gly Val 345

Ser Glu Ser Gln Pro Val Val Arg Lys Arg Thr Tyr Arg Lys Leu Cys 365 360 355

Ala Ile His His Leu Leu Gly Arg Ala Pro Glu Ile Ala Cys Ser Ala 375 370

Cys Asp Leu Val Asn Val Asp Leu Asp Asp Cys Val Ser Glu Gln 395 390 385

<210> 10

<211> 969

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1)..(966)

<220>

<221> misc feature

<222> (943)..(944)

<223> minor splice site

<400> 10

atg gag ctg gtc ggg tgg ctg gtg gac cgg ggc atc acc tcc gag aag Met Glu Leu Val Gly Trp Leu Val Asp Arg Gly Ile Thr Ser Glu Lys 15 10

cag tgg atc cag gag gac cag gcc tcg tac atc tcc ttc aac gcc gct Gln Trp Ile Gln Glu Asp Gln Ala Ser Tyr Ile Ser Phe Asn Ala Ala 25 20

tcc aac tcg cgg tcc cag atc aag gcc gct ctg gac aat gcc ggc aag 144

Ser	Asn	Ser 35		Ser	Gln	Ile	Lys 40		Ala	Leu	Asp	Asn 45	Ala	Gly	Lys	
		Ala								tac Tyr						192
_										tac Tyr 75						240
			-							gtc Val						288
										tgg Trp						336
	_									atc Ile						384
										aac Asn						432
										gag Glu 155						480
_	-									ggc						528
	-	caa Gln	-	-						atc Ile						576
	-					Asn				gtg Val						624
			-							gac Asp						672
gaa	ctc	acc	cgc	cgt	ctg	gag	cat	gac	ttt	ggc	aag	gtg	aca	aag	cag	720

Glu 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gl.y 235	Lys	Val	Thr	Lys	Gln 240	
~~~	atc	222	aaa	ttc	ttc	cac	taa	aca	сап	gat	cac	ata	acc	gag	ata	768
														Glu 255	Val	
														ece Pro		816
	-													tca Ser		864
														ttt Phe		912
-				_										gga Gly		960
	ctc Leu	tga														969
<213 <212	0> 11 l> 32 2> PF 8> AF	22 RT	•													
	)> 11	-					_						_		_	
Met 1	Glu	Leu	Val	Gly 5	Trp	Leu	Val	Asp	Arg 10	Gly	Ile	Thr	Ser	Glu 15	Lys	
Gln	Trp	Ile	Gln 20	Glu	Asp	Gln	Ala	Ser 25	Туr	Ile	Ser	Phe	Asn 30	Ala	Ala	
Ser	Asn	Ser 35	Arg	Ser	Gln	Ile	Lys 40	Ala	Ala	Leu	Asp	Asn 45	Ala	Gly	Lys	
Ile	Met 50	Ala	Leu	Thr	Lys	Ser 55	Ala	Pro	Asp	Tyr	Leu 60	Val	Gly	Pro	Ala	
Pro 65	Pro	Ala	Asp	Ile	Lys 70	Thr	Asn	Arg	Ile	Туг 75	Arg	Ile	Leu	Glu	Leu 80	

Asn	Gly	Туr	Glu	Pro 85	۸la	Туr	Ala	Gly	Ser 90	Val	Phe	Leu	Gly	Trp 95	Ala
Gln	Lys	Arg	Phe 100	Gly	Lys	Arg	Asn	Thr 105	Ile	Trp	Leu	Phe	Gly 110	Pro	Ala
Thr	Thr	Gly 115	Lys	Thr	Asn	Ile	Ala 120	Glu	Ala	Ile	Ala	His 125	Ala	Val	Pro
. Phe	Tyr 130	Gly	Cys	Val	Asn	Тгр 135	Thr	Asn	Glu	Asn	Phe 140	Pro	Phe	Asn	Asp
Cys 145	Val	Asp	Lys	Met	Val 150	Ile	Trp	Trp	Glu	Glu 155	Gly	Lys	Met	Thr	Ala 160
Lys	Val	Val	Glu	Ser 165	Ala	Lys	Ala	Ile	Leu 170	Gly	Gly	Ser	Lys	Val 175	Arg
Val	Asp	Gln	Lys 180	Cys	Lys	Ser	Ser	Ala 185	Gln	Ile	Asp	Pro	Thr 190	Pro	Val
Ile	Val	Thr 195	Ser	Asn	Thr	Asn	Met 200	Cys	Ala	Val	Ile	Asp 205	Gly	Asn	Ser
Thr	Thr 210	Phe	Glu	His	Gln	Gln 215	Pro	Leu	Gln	Asp	Arg 220	Met	Phe	Lys	Phe
Glu 225	Leu	Thr	Arg	Arg	Leu 230	Glu	His	Asp	Phe	Gly 235	Lys	Val	Thr	Lys	Gln 240
Glu	Val	Lys	Glu	Phe 245	Phe	Arg	Trp	Ala	Gln 250	Asp	His	Val	Thr	Glu 255	Val
Ala	His	Glu	Phe 260	Туr	Val	Arg	Lys	Gly 265	Gly	Ala	Asn	Lys	Arg 270	Pro	Ala
Pro	Asp	Asp 275	Ala	Asp	Lys	Ser	Glu 280	Pro	Lys	Arg	Ala	Cys 285	Pro	Ser	Val
Ala	Asp 290	Pro	Ser	Thr	Ser	Asp 295	Ala	Glu	Gly	Ala	Pro 300	Val	Asp	Phe	Ala
Asp 305	Arg	Tyr	Gly	Cys	Arg 310	Trp	Leu	Ser	Ser	Arg <b>3</b> 15	Leu	Ala	Arg	Gly	Gln 320

Pro Leu

<21	0> 1 1> 2	211														
	2> D 3> A	NA AV-1														
	1> C	DS 1)	(220	8)												
<40	0> 1	2														
_	-				Tyr					Leu			aac Asn			48
													ccg Pro 30			96
	_												gtg Val			144
													Gly			192
_				-	-								gcc Ala			240
													aac Asn			288
_							Leu						ttt Phe 110			336
													ctc Leu			384
		-	-										aag Lys			432
cca	qta	gag	cag	tca	cca	caa	gag	cca	gac	tcc	tcc	tca	qqc	atc	ggc	480

Pro 145		Glu	Gln	Ser	Pro 150	Gln	Glu	Pro	Asp	Ser 155	Ser	Ser	Gly	Ile	<b>Gly</b> 160	
_		Glà														528
	-	tca Ser														576
_		ccc Pro 195		-												624
_		atg Met	-	-												672
	_	aat Asn														720
		agc Ser		•			-	_								768
	_	caa Gln														816
		ggc Gly 275														864
His	Cys 290	cac His	Phe	Ser	Pro	Arg 295	Asp	Trp	Gln	Arg	Leu 300	Ile	Asn	Asn	Asn	912
Trp 305	Gly	ttc Phe	Arg	Pro	Lys 310	Arg	Leu	Asn	Phe	Lys 315	Leu	Phe	Asn	Ile	Gln 320	960
-	_	gag Glu														1008
ctt	acc	agc	acg	gtt	caa	gtc	ttc	tcg	gac	tcg	gag	tac	cag	ctt	ccg	1056

•			_													
Leu	Thr	Ser	Thr 340	Val	Gln	Val	Phe	ser 345	Asp	Ser	Glu	Tyr	<b>Gl</b> n <b>3</b> 50	Leu	Pro	
tac Tyr	gtc Val	ctc Leu 355	ggc Gly	tct Ser	gcg Ala	cac His	cag Gln 360	ggc Gly	tgc Cys	ctc Leu	cct Pro	ccg Pro 365	ttc Phe	ccg Pro	gcg Ala	1104
gac Asp	gtg Val 370	ttc Phe	atg Met	att Ile	ccg Pro	caa Gln 375	tac Tyr	ggc Gly	tac Tyr	ctg Leu	acg Thr 380	ctc Leu	aac Asn	aat Asn	ggc ggc	1152
agc Ser 385	caa Gln	gcc Ala	gtg Val	gga Gly	cgt Arg 390	tca Ser	tcc Ser	ttt Phe	tac Tyr	tgc Cys 395	ctg Leu	gaa Glu	tat Tyr	ttc Phe	cct Pro 400	1200
tct Ser	cag Gln	atg Met	ctg Leu	aga Arg 405	acg Thr	ggc Gly	aac Asn	aac Asn	ttt Phe 410	acc Thr	ttc Phe	agc Ser	tac Tyr	acc Thr 415	ttt Phe	1248
gag Glu	gaa Glu	gtg Val	cct Pro 420	ttc Phe	cac His	agc Ser	agc Ser	tac Tyr 425	gcg Ala	cac His	agc Ser	cag Gln	agc Ser 430	ctg Leu	gac Asp	1296
cgg Arg	ctg Leu	atg Met 435	aat Asn	cct Pro	ctc Leu	atc Ile	gac Asp 440	caa Gln	tac Tyr	ctg Leu	tat Tyr	tac Tyr 445	ctg Leu	aac Asn	aga Arg	1344
act Thr	caa Gln 450	aat Asn	cag Gln	tcc Ser	gga Gly	agt Ser 455	gcc Ala	caa Gln	aac Asn	aag Lys	gac Asp 460	ttg Leu	ctg Leu	ttt Phe	agc Ser	1392
cgt Arg 465	ggg ggg	tct Ser	cca Pro	gct Ala	ggc Gly 470	atg Met	tct Ser	gtt Val	cag Gln	ccc Pro 475	aaa Lys	aac Asn	tgg Trp	cta Leu	cct Pro 480	1440
gga Gly	ccc Pro	tgt Cys	tat Tyr	cgg Arg 485	cag Gln	cag Gln	cgc Arg	gtt Val	tct Ser 490	aaa Lys	aca Thr	aaa Lys	aca Thr	gac Asp 495	aac Asn	1488
aac Asn	aac Asn	agc Ser	aat Asn 500	ttt Phe	acc Thr	tgg Trp	act Thr	ggt Gly 505	gct Ala	tca Ser	aaa Lys	tat Tyr	aac Asn 510	ctc Leu	aat Asn	1536
ggg	cgt Arg	gaa Glu 515	tcc Ser	atc Ile	atc Ile	aac Asn	cct Pro 520	ggc Gly	act Thr	gct Ala	atg Met	gcc Ala 525	tca Ser	cac His	aaa Lys	1584
gac	gac	gạa	gac	aag	ttc	ttt	ccc	atg	agc	ggt	gtc	atg	att	ttt	gga	1632

Asp	Asp 530	Glu	Asp	Lys	Phe	Phe 535	Pro	Met	Ser	Gly	Val 540	Met	Ile	Phe	Gly	
aaa Lys 545	gag Glu	agc Ser	gcc Ala	gga Gly	gct Ala 550	tca Ser	aac Asn	act Thr	gca Ala	ttg Leu 555	gac Asp	aat Asn	gtc Val	atg Met	att Ile 560	1680
					att Ile											1728
					gtc Val											1776
					gct Ala											1824
					ctg Leu											1872
				Phe	cac His 630											1920
					cag Gln											1968
					ttt Phe											2016
caa Gln	tac Tyr	tcc Ser 675	aca Thr	gga Gly	caa Gln	gtg Val	agt Ser 680	gtg <b>V</b> al	gaa Glu	att Ile	gaa Glu	tgg Trp 685	gag Glu	ctg Leu	cag Gln	2064
					cgc Arg											2112
					aac Asn 710											2160
tat	act	gag	cct	cgc	ccc	att	ggc	acc	cgt	tac	ctt	acc	cgt	ccc	ctg	2208

37 .

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

taa 2211

<210> 13

<211> 736

<212> PRT

<213> AAV-1

<400> 13

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser

1 5 10 15

Glu Gly Ile Arg Glu Trp Trp Asp Leu Lys Pro Gly Ala Pro Lys Pro 20 25 30

Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro 35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro . 50 55 60

Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp 65 70 75 80

Gln Gln Leu Lys Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala 85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro 115 120 125

Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg 130 135 140

Pro Val Glu Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Ile Gly
145 150 155 160

Lys Thr Gly Gln Gln Pro Ala Lys Lys Arg Leu Asn Phe Gly Gln Thr 165 170 175

Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro 180 185 190

Ala Thr Pro Ala Ala Val Gly Pro Thr Thr Met Ala Ser Gly Gly Gly
195 . 200 205

Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ala 210 215 220

Ser Gly Asn Trp His Cys Asp Ser Thr Trp Leu Gly Asp Arg Val Ile 225 230 235 240

Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu 245 250 255

Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly Ala Ser Asn Asp Asn His 260 265 270

Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe 275 280 285

His Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn 290 295 300

Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe Lys Leu Phe Asn Ile Gln 305 310 315 320

Val Lys Glu Val Thr Thr Asn Asp Gly Val Thr Thr Ile Ala Asn Asn 325 330 335

Leu Thr Ser Thr Val Gln Val Phe Ser Asp Ser Glu Tyr Gln Leu Pro 340 345 350

Tyr Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala 355 360 365

Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly 370 375 380

Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro 385 390 395 400

Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe
405 410 415

Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp 420 425 430

Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg
435 440 445

Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser 450 455 460

- Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro 465 470 475 480
- Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn 485 490 495
- Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn 500 505 510
- Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys 515 520 525
- Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly 530 535 540
- Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile 545 550 555 560
- Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg 565 570 575
- Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala 580 585 590
- Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln
  595 600 605
- Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His 610 615 620
- Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu 625 635 640
- Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys Asn Thr Pro Val Pro Ala 645 650 655
- Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys Phe Ala Ser Phe Ile Thr 660 665 670
- Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln 675 680 685
- Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Val Gln Tyr Thr Ser Asn 690 695 700

Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr Val Asp Asn Asn Gly Leu 705 710 715. 720

Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Pro Leu
725 730 735

<210> 14

<211> 1800

<212> DNA

<213> AAV-1

<220>

<221> CDS

<222> (1) .. (1797)

<400> 14

acg gct cct gga aag aaa cgt ccg gta gag cag tcg cca caa gag cca 48
Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro
1 5 10 15

gac tcc tcc tcg ggc atc ggc aag aca ggc cag cag ccc gct aaa aag 96
Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys
20 25 30

aga ctc aat ttt ggt cag act ggc gac tca gag tca gtc ccc gat cca 144
Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro
35 40 45

caa cct ctc gga gaa cct cca gca acc ccc gct gct gtg gga cct act 192
Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr
50 55 60

aca atg gct tca ggc ggt ggc gca cca atg gca gac aat aac gaa ggc 240
Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly
65 70 75 80

gcc gac gga gtg ggt aat gcc tca gga aat tgg cat tgc gat tcc aca 288
Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr
85 90 95

tgg ctg ggc gac aga gtc atc acc acc agc acc cgc acc tgg gcc ttg 336
Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu
100 105 110

ccc acc tac aat aac cac ctc tac aag caa atc tcc agt gct tca acg 384
Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr
115 120 125

ggg Gly	gcc Ala 130	agc Ser	aac Asn	gac Asp	aac Asn	cac His 135	tac Tyr	ttc. Phe	ggc Gly	tac Tyr	agc. Ser 140	acc Thr	ccc Pro	tgg. Trp	ggg Gly	432
tat Tyr 145	ttt Phe	gat Asp	ttc Phe	aac Asn	aga Arg 150	ttc Phe	cac His	tgc Cys	cac His	ttt Phe 155	tca Ser	cca Pro	cgt Arg	gac Asp	tgg Trp 160	480
cag Gln	cga Arg	ctc Leu	atc Ile	aac Asn 165	aac Asn	aat Asn	tgg Trp	gga Gly	ttc Phe 170	cgg Arg	ccc Pro	aag Lys	aga Arg	ctc Leu 175	aac Asn	528
ttc Phe	aaa Lys	ctc Leu	ttc Phe 180	aac Asn	atc Ile	caa Gln	gtc Val	aag Lys 185	gag Glu	gtc Val	acg Thr	acg Thr	aat Asn 190	gat Asp	Gly Gly	576
gtc Val	aca Thr	acc Thr 195	atc Ile	gct Ala	aat Asn	aac Asn	ctt Leu 200	acc Thr	agc Ser	acg Thr	gtt Val	caa Gln 205	gtc Val	ttc Phe	tcg Ser	624
gac Asp	tcg Ser 210	gag Glu	tac Tyr	cag Gln	ctt Leu	ccg Pro 215	tac Tyr	gtc Val	ctc Leu	ggc Gly	tct Ser 220	gcg Ala	cac His	cag Gln	ggc Gly	672
tgc Cys 225	ctc Leu	cct Pro	ccg Pro	ttc Phe	ccg Pro 230	gcg Ala	gac Asp	gtg Val	ttc Phe	atg Met 235	att Ile	ccg Pro	caa Gln	tac Tyr	ggc Gly 240	720
tac Tyr	ctg Leu	acg Thr	ctc Leu	aac Asn 245	aat Asn	ggc Gly	agc Ser	caa Gln	gcc Ala 250	gtg Val	gga Gly	cgt Arg	tca Ser	ser 255	ttt Phe	768
Tyr	Cys	Leu	Glu 260	Tyr	ttc Phe	Pro	Ser	Gln 265	Met	Leu	Arg	Thr	Gly 270	Asn	Asn	816
Phe	Thr	Phe 275	Ser	Tyr	acc Thr	Phe	Glu 280	Glu	Val	Pro	Phe	His 285	Ser	Ser	Tyr	864
gcg Ala	cac His 290	agc Ser	cag Gln	agc Ser	ctg Leu	gac Asp 295	cgg Arg	ctg Leu	atg Met	aat Asn	cct Pro 300	ctc Leu	atc Ile	gac Asp	caa Gln	912
tac Tyr 305	ctg Leu	tat Tyr	tac Tyr	ctg Leu	aac Asn 310	aga Arg	act Thr	çaa Gln	aat Asn	cag Gln 315	tcc Ser	gga Gly	agt Ser	gcc Ala	caa Gln 320	960

					Phe			G1 y								1008
								ccc Pro 345								1056
								aac Asn								1104
-								cgt Arg								1152
	-	_	-					gac Asp								1200
-		-	-					gag Glu								1248
-	-	_		-				gac Asp 425								1296
			-		-			G] À aaa								1344
Ser	Ser 450	Ser	Thr	Asp	Pro	Ala 455	Thr	gga Gly	Asp	Val	His 460	Ala	Met	Gly	Ala	1392
Leu 465	Pro	Gly	Met	Val	Trp 470	Gln	Asp	aga Arg	Asp	Val 475	Tyr	Leu	Gln	Gly	Pro 480	1440
		_						gat Asp								1488
	-							aac Asn 505				Gln				1536

aaa aac acg cct gtt cct gcg aat cct ccg gcg gag ttt tca gct aca Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 525 520 515 aag ttt gct tca ttc atc acc caa tac tcc aca gga caa gtg agt gtg Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 530 535 540 gaa att gaa tgg gag ctg cag aaa gaa aac agc aag cgc tgg aat ccc 1680 Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 555 545 550 gaa gtg cag tac aca tcc aat tat gca aaa tct gcc aac gtt gat ttt 1728 Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe 570 565 act gtg gac aac aat gga ctt tat act gag cct cgc ccc att ggc acc 1776 Thr Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr 585 580 1800 cgt tac ctt acc cgt ccc ctg taa Arg Tyr Leu Thr Arg Pro Leu 595 <210> 15 <211> 599 <212> PRT <213> AAV-1 <400> 15 Thr Ala Pro Gly Lys Lys Arg Pro Val Glu Gln Ser Pro Gln Glu Pro 10 Asp Ser Ser Ser Gly Ile Gly Lys Thr Gly Gln Gln Pro Ala Lys Lys 25 Arg Leu Asn Phe Gly Gln Thr Gly Asp Ser Glu Ser Val Pro Asp Pro 45 40 35 Gln Pro Leu Gly Glu Pro Pro Ala Thr Pro Ala Ala Val Gly Pro Thr 50 55 Thr Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly 65 70 75

Ala Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr

85 90 95

Trp Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu 100. 105 110

- Pro Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr 115 120 125
- Gly Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
  130 135 140
- Tyr Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp 145 150 155 160
- Gln Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn 165 170 175
- Phe Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly
  180 185 190
- Val Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser 195 200 205
- Asp Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly
  210 215 220
- Cys Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly
  225 230 235 240
- Tyr Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe 245 250 255
- Tyr Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn 260 265 270
- Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr 275 280 285
- Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln 290 295 300
- Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln 305 310 315 320
- Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val 325 330 335
- Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val

340

345

350

Ser Lys Thr Lys Thr Asp Asn Asn Ser Asn Phe Thr Trp Thr Gly 360

Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly 375

Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met 390 395

Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr

Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr 425 420

Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln 440

Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala 455

Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro 470 475

Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro 490 485

Leu Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile . 500

Lys Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr 520

Lys Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val 535

Glu Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro 545 550

Glu Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe 565 570

Thr Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr **58**5 580

Arg Tyr Leu Thr Arg Pro Leu

<21	0> 1	6														
<21	1> 1	605														
<21	2> D	NA														
<21	3> A	AV-1														
<22	0>															
<22	1> C	DS														
<22	2> (	1)	(160	2)												
	0> 1															
_	-							atg								48
Met	Ala	Ser	Gly			Ala	Pro	Met		Asp	Asn	Asn	GLu		Ala	
1				5					10		٠			15		
													<b>.</b>		<b>.</b>	0.0
-					_			aat								96
Asp	Gly	Val	_	Asn	Ala	ser	GLY	Asn	тгр	HIS	Cys	Asp		The	тър	
			20					25					30			
ata.	<b>77</b>	<b>42</b> C	202	atc	atc	200	300	agc	acc	cac	acc	taa	acc	tta	ccc	144
_		_	-	_				Ser								
ьец	GLY	жар 35	ALG	Val	116	1111	40	SCL	1111	ALY	1111	45	ALG	БСи	110	
		33					10					10				
acc	tac	aat	aac	cac	ctc	tac	aag	caa	atc	tcc	agt	act	tca	acq	aaa	192
							-	Gln			_	-		_		
	50					55	•				60				-	
gcc	agc	aac	gac	aac	cac	tac	ttc	ggc	tac	agc	acc	ccc	tgg	ggg	tat	240
Ala	Ser	Asn	Asp	Asn	His	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr	
65					70					75					80	
ttt	gat	ttc	aac	aga	ttc	cac	tgc	cac	ttt	tca	cca	cgt	gac	tgg	cag	288
Phe	Asp	Phe	Asn	Arg	Phe	His	Cys	His	Phe	Ser	Pro	Arg	Asp	Trp	Gln	
				85					90					95		
cga	ctc	atc	aac	aac	aat	tgg	gga	ttc	cgg	ccc	aag	aga	ctc	aac	ttc	336
Arg	Leu	Ile	Asn	Asn	Asn	$\operatorname{Trp}$	Gly	Phe	Arg	Pro	Lys	Arg	Leu	Asn	Phe	
			100					105					110			
				•		-	_	gag	-	_	_		-			384
Lys	Leu	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Thr	Asn	Asp	Gly	Val	
		115					120					125				
			-					agc	-	-		-				432
Thr		Ile	Ala	Asn	Asn		Thr	Ser	Thr	Val		Val	Phe	Ser	Asp	
	130					125					140					

tcg	gag	tac	cag	ctt	ccg	tac	gtc	ctc	ggc	tct	gcg	cac	cag	ggc Gl v	tgc.	480
Ser 145	Glu	Туr	Gln	Leu	Pro 150	Tyr	Val	Leu	GIŸ	155	AIa	1113	Gln	O <sub>2</sub> y	160	
ctc Leu	cct Pro	ccg Pro	ttc Phe	ccg Pro	gcg Ala	gac Asp	gtg Val	ttc Phe	atg Met	att Ile	ccg Pro	caa Gln	tac Tyr	ggc Gly	tac Tyr	528
				165					170					175		576
ctg Leu	acg Thr	ctc Leu	aac Asn 180	aat Asn	ggc Gly	agc Ser	caa Gln	gcc Ala 185	yal Val	gga Gly	Arg	Ser	tcc Ser 190	Phe	Tyr	370
tac	cta	gaa		ttc	cct	tct	cag		ctg	aga	acg	ggc	aac	aac	<b>t</b> tt	624
Cys	Leu	Glu 195	Tyr	Phe	Pro	Ser	Gln 200	Met	Leu	Arg	Thr	Gly 205	Asn	Asn	Phe	
acc	ttc	agc	tac	acc	ttt	gag	gaa	gtg Val	cct Pro	ttc Phe	cac His	agc Ser	agc Ser	tac Tyr	gcg Ala	672
Thr	210	Ser	tàr	1111	riie	215	oru	742			220			-		
cac His	agc Ser	cag Gln	agc Ser	ctg Leu	gac Asp	cgg Arg	ctg Leu	atg Met	aat Asn	Pro	ctc Leu	atc Ile	gac Asp	caa Gln	Tyr.	720
225					230	3.C.T	C22	aat	can	235	gga	agt	gcc	caa	240 aac	768
Leu	Tyr	Tyr	Leu	Asn 245	Arg	Thr	Gln	Asn	Gln 250	Ser	Gly	Ser	Ála	Gln 255	Asn	
aag	gac	ttg	ctg	ttt	agc	cgt	ggg	tct	cca	gct Ala	ggc Gl v	atg Met	tct Ser	gtt Val	cag Gln	816
Lys	Asp	Leu	260	rne	Ser	ALG	GIY	265	110	,,,,	0_,		270			
ccc Pro	aaa Lys	aac Asn	tgg Trp	cta Leu	cct Pro	gga Gly	ccc Pro	tgt Cys	tat Tyr	cgg Arg	cag Gln	Gln	cgc Arg	gtt Val	tct Ser	864
		275		<b></b>	226	220	280	anc	aat	EEE	acc	285 tag	act	ggt	gct	912
aaa Lys	Thr 290	Lys	Thr	Asp	Asn	Asn 295	Asn	Ser	Asn	Phe	Thr 300	Trp	Thr	Gly	Ala	
tca	aaa	tat	aac	ctc	aat	ggg	cgt	gaa	tcc	atc	atc	aac	cct	ggc	act	960
Ser 305	Lys	Tyr	Asn	Leu	Asn 310	Gly	Arg	Glu	Ser	11e 315	Ile	Asn	Pro	GTÀ	320	
gct Ala	atg Met	gcc Ala	tca Ser	cac His	aaa Lvs	gac Asp	gac Asp	gaa Glu	gac Asp	aag Lys	ttc Phe	ttt Phe	ccc Pro	atg Met	agc Ser	1008
				325	-	•	-		330					335		

				Phe	gga Gly						1056
					att Ile						1104
		Ala			aga Arg						1152
_	-		-		gcg Ala 390						1200
					caa Gln						1248
					cac His						1296
-					ctc Leu						1344
	-		-		gcg Ala						1392
	_				acc Thr 470						1440
	-	-			cag Gln						1488
	_				aat Asn				Val		1536
	_				ctt Leu	Tyr	-				1584

1605

tac ctt acc cgt ccc ctg taa Tyr Leu Thr Arg Pro Leu 530

<210> 17
<211> 534
<212> PRT
<213> AAV-1

<400> 17

Met Ala Ser Gly Gly Gly Ala Pro Met Ala Asp Asn Asn Glu Gly Ala 1 5 10 15

Asp Gly Val Gly Asn Ala Ser Gly Asn Trp His Cys Asp Ser Thr Trp 20 25 30

Leu Gly Asp Arg Val Ile Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro 35 40 45

Thr Tyr Asn Asn His Leu Tyr Lys Gln Ile Ser Ser Ala Ser Thr Gly 50 55 60

Ala Ser Asn Asp Asn His Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr 65 70 75 80

Phe Asp Phe Asn Arg Phe His Cys His Phe Ser Pro Arg Asp Trp Gln 85 90 95

Arg Leu Ile Asn Asn Asn Trp Gly Phe Arg Pro Lys Arg Leu Asn Phe 100 105 110

Lys Leu Phe Asn Ile Gln Val Lys Glu Val Thr Thr Asn Asp Gly Val 115 120 125

Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Ser Asp 130 135 140

Ser Glu Tyr Gln Leu Pro Tyr Val Leu Gly Ser Ala His Gln Gly Cys 145 150 155 160

Leu Pro Pro Phe Pro Ala Asp Val Phe Met Ile Pro Gln Tyr Gly Tyr 165 170 175

Leu Thr Leu Asn Asn Gly Ser Gln Ala Val Gly Arg Ser Ser Phe Tyr 180 185 190

Cys Leu Glu Tyr Phe Pro Ser Gln Met Leu Arg Thr Gly Asn Asn Phe Thr Phe Ser Tyr Thr Phe Glu Glu Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr Gln Asn Gln Ser Gly Ser Ala Gln Asn Lys Asp Leu Leu Phe Ser Arg Gly Ser Pro Ala Gly Met Ser Val Gln Pro Lys Asn Trp Leu Pro Gly Pro Cys Tyr Arg Gln Gln Arg Val Ser Lys Thr Lys Thr Asp Asn Asn Asn Ser Asn Phe Thr Trp Thr Gly Ala Ser Lys Tyr Asn Leu Asn Gly Arg Glu Ser Ile Ile Asn Pro Gly Thr Ala Met Ala Ser His Lys Asp Asp Glu Asp Lys Phe Phe Pro Met Ser Gly Val Met Ile Phe Gly Lys Glu Ser Ala Gly Ala Ser Asn Thr Ala Leu Asp Asn Val Met Ile Thr Asp Glu Glu Glu Ile Lys Ala Thr Asn Pro Val Ala Thr Glu Arg Phe Gly Thr Val Ala Val Asn Phe Gln Ser Ser Ser Thr Asp Pro Ala Thr Gly Asp Val His Ala Met Gly Ala Leu Pro Gly Met Val Trp Gln Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His Thr Asp Gly His Phe His Pro Ser Pro Leu 

Met Gly Gly Phe Gly Leu Lys Asn Pro Pro Pro Gln Ile Leu Ile Lys
435 440 445

Asn Thr Pro Val Pro Ala Asn Pro Pro Ala Glu Phe Ser Ala Thr Lys 450 455 460

Phe Ala Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Ser Val Glu 465 470 475 480

Ile Glu Trp Glu Leu Gln Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu 485 490 495

Val Gln Tyr Thr Ser Asn Tyr Ala Lys Ser Ala Asn Val Asp Phe Thr 500 505 510

Val Asp Asn Asn Gly Leu Tyr Thr Glu Pro Arg Pro Ile Gly Thr Arg 515 520 525

Tyr Leu Thr Arg Pro Leu 530

<210> 18

<211> 4681

<212> DNA

<213> aav-2

<400> 18

ttggccacte cetetetgeg egetegeteg eteactgagg eegggegaee aaaggtegee 60 egacgecegg getttgeeeg ggeggeetea gtgagegage gagegegaag agagggagtg 120 gecaacteea teactagggg tteetggagg ggtggagteg tgacgtgaat taegteatag 180 ggttagggag gteetgtatt agagggeaeg tgagtgttt gegacatttt gegacaccat 240 gtggtcacge tgggtatta agecegagtg ageacgeagg gtetecattt tgaageggga 300 ggtttgaacg egeageegee atgeegggt tttaegagat tgtgattaag gteeceageg 360 accttgacgg geatetgeee ggeattetg acagettgt gaactgggtg geegagaagg 420 aatgggagtt geegeagat tetgacatgg atetgaatet gattgageag geaceeetga 480 eegtggeega gaagetgeag egegactte tgaeggaatg geegegtgt agtaaggeee 540 eggaggeeet tttettgtg caatttgaga agggagagag etaetteeae atgeaegtge 600 tegtggaaac caeeggggtg aaateeatgg tttteggage ttteetgagt cagattegeg 720 aaaaaactgat teagagaatt taeegeggga tegageegae ttttgecaaac tggttegegg 720

tcacaaagac cagaaatggc gccggaggcg ggaacaaggt ggtggatgag tgctacatcc 780 ccaattactt gctccccaaa acccagcctg agctccagtg ggcgtggact aatatggaac 840 agtatttaag cgcctgtttg aatctcacgg agcgtaaacg gttggtggcg cagcatctga 900 cgcacgtgtc gcagacgcag gagcagaaca aagagaatca gaatcccaat totgatgcgc 960 cggtgatcag atcaaaaact tcagccaggt acatggagct ggtcgggtgg ctcgtggaca 1020 aggggattac ctcggagaag cagtggatcc aggaggacca ggcctcatac atctccttca 1080 atgcggcctc caactcgcgg tcccaaatca aggctgcctt ggacaatgcg ggaaagatta 1140 tgagcctgac taaaaccgcc cccgactacc tggtgggcca gcagcccgtg gaggacattt 1200 ccagcaatcg gatttataaa attttggaac taaacgggta cgatccccaa tatgcggctt 1260 ccgtctttct gggatgggcc acgaaaaagt tcggcaagag gaacaccatc tggctgtttg 1320 ggcctgcaac taccgggaag accaacatcg cggaggccat agcccacact gtgcccttct 1380 acgggtgcgt aaactggacc aatgagaact ttcccttcaa cgactgtgtc gacaagatgg 1440 tgatctggtg ggaggagggg aagatgaccg ccaaggtcgt ggagtcggcc aaagccattc 1500 tcggaggaag caaggtgcgc gtggaccaga aatgcaagtc ctcggcccag atagacccga 1560 ctcccgtgat cgtcacctcc aacaccaaca tgtgcgccgt gattgacggg aactcaacga 1620 ccttcgaaca ccagcagecg ttgcaagace ggatgt@caa atttgaactc acccgccgte 1680 tggatcatga ctttgggaag gtcaccaagc aggaagtcaa agactttttc cggtgggcaa 1740 aggatcacgt ggttgaggtg gagcatgaat tctacgtcaa aaagggtgga gccaagaaaa 1800 gacccgcccc cagtgacgca gatataagtg agcccaaacg ggtgcgcgag tcagttgcgc 1860 agccatcgac gtcagacgcg gaagcttcga tcaactacgc agacaggtac caaaacaaat 1920 gttctcgtca cgtgggcatg aatctgatgc tgtttccctg cagacaatgc gagagaatga 1980 atcagaattc aaatatctgc ttcactcacg gacagaaaga ctgtttagag tgctttcccg 2040 tgtcagaatc tcaaccggtt tctgtcgtca aaaaggcgta tcagaaactg tgctacattc 2100 atcatatcat gggaaaggtg ccagacqctt gcactgcctg cgatctggtc aatgtggatt 2160

tggatgactg catctttgaa caataaatga tttaaatcag gtatggctgc cgatggttat 2220 cttccagatt ggctcgagga cactctctct gaaggaataa gacagtggtg gaagctcaaa 2280 cctggcccac caccaccaaa gcccgcagag cggcataagg acgacagcag gggtcttgtg 2340 ctteetgggt acaagtacet eggaceette aacggacteg acaagggaga geeggteaac 2400 gaggcagacg ccgcggccct cgagcacgac anagcctacg accggcagct cgacagcgga 2460 gacaacccgt acctcaagta caaccacgcc gacgcggagt ttcaggagcg ccttaaagaa 2520 gatacgtctt ttgggggcaa cctcggacga gcagtcttcc aggcgaaaaa gagggttctt 2580 gaaceteteg geetggttga ggaacetgtt aagaeggete egggaaaaaa gaggeeggta 2640 gagcactete etgtggagee agacteetee tegggaaceg gaaageeggg ceagcageet 2700 gcaagaaaaa gattgaattt tggtcagact ggagacgcag actcagtacc tgacccccag 2760 ceteteggae agecaceage agececetet ggtetgggaa etaataegat ggetacagge 2820 agtggcgcac caatggcaga caataacgag ggcgccgacg gagtgggtaa ttcctccgga 2880 aattggcatt gcgattccac atggatgggc gacagagtca tcaccaccag cacccgaacc 2940 tgggccctgc ccacctacaa caaccacctc tacaaacaaa tttccagcca atcaggagcc 3000 togaacgaca atcactactt tygctacago acceettggg ggtattttga ettcaacaga 3060 ttocactgoe acttttcace aegtgactgg caaagactea teaacaacaa etggggatte 3120 cgacccaaga gactcaactt caacctcttt aacattcaag tcaaagaggt cacgcagaat 3180 gacggtacga cgacgattgc caataacctt accagcacgg ttcaggtgtt tactgactcg 3240 gagtaccage tecegtacgt ecteggeteg gegeateaag gatgeeteee geegtteeea 3300 gcagacgtct tcatggtgcc acagtatgga tacctcaccc tgaacaacgg gagtcaggca 3360 graggacget etteattta etgeetggag taettteett eteagatget gegtaeegga 3420 aacaacttta cetteageta caettttgag gaegtteett tecacageag etaegeteae 3480 agccagagte tggaccgtet catgaateet eteategace agtacetgta ttaettgage 3540 agaacaaaca ctccaagtgg aaccaccacg cagtcaaggc ttcagttttc tcaggcccca 3600

gccagtgaca ttcgggacca gtctaggaac tggcttcctg gaccctgtta ccgccagcag 3660 cgagtatgaa agacatctgc ggataacaac aacagtgaat actcgtggac tggagctacc 3720 aagtaccace tcaatggcag agactetetg gtgaateegg ggeregeeat ggcaageeac 3780 aaggacgatg aagaaaagtt ttttcctcag ageggggttc tcatctttgg gaagcaagge 3840 tcagagaaaa caaatgtgaa cattgaaaag gtcatgatta cagacgaaga ggaaatccca 3900 acaaccaatc ccgtggctac ggagcagtat ggttctgtat ctaccaacct ccagagaggc 3960 aacagacaag cagctaccgc agatgtcaac acacaaggcg ttcttccagg catggtctgg 4020 caggacagag atgtgtacct tcaggggccc atctgggcaa agattccaca cacggacgga 4080 cattttcacc cctctcccct catgggtgga ttcggactta aacaccctcc tccacagatt 4140 ctcatcaaga acaccccggt acctgcgaat ccttcgacca ccttcagtgc ggcaaagttt 4200 getteettea teacacagta etecaeggga caeggteage gtggagateg agtgggaget 4260 gcagaacgaa aacagcaaac gctggaatcc cgaaattcag tacacttcca actacaacaa 4320 gtotgttaat ogtggacttt accgtggata otaatggogt gtattoagag ootogooca 4380 ttggcaccag atacctgact cgtaatctgt aattgcttgt taatcaataa accgtttaat 4440 togtttcagt tgaactttgg tototgogra tttotttott atotagttto catggotacg 4500 tagataagta gcatggcggg ttaatcabta actacaagga acccctagtg atggagttgg 4560 ccactccctc tetgegeget egetegetea etgaggeegg gegaccaaag gtegecegae 4620 gcccgggctt tgccccggcg gcctcagtga gcgagcgagc gcgcagagag ggagtgggca 4680 4681 a

<sup>&</sup>lt;210> 19

<sup>&</sup>lt;211> 4683

<sup>&</sup>lt;212> DNA

<sup>&</sup>lt;213> aav-6

<sup>&</sup>lt;400> 19

ttggccactc cctctctgcg cgctcgctcg ctcactgagg ccgggcgacc aaaggtcgcc 60

cgacgcccgg gctttgcccg ggcggcctca gtgagcgagc gagcgcgcag agagggagtg 120 gccaacteca teactagggg tteetggagg ggtggagteg tgaegtgaat taegteatag 180 ggttagggag gtcctgtatt agaggtcacg tgagtgtttt gcgacatttt gcgacaccat 240 gtggtcacgc tgggtattta agcccgagtg agcacgcagg gtctccattt tgaagcggga 300 ggtttgaacg cgcagcgcca tgccggggtt ttacgagatt gtgattaagg tccccagcga 360 ccttgacgag catctgcccg gcatttctga cagctttgtg aactgggtgg ccgagaagga 420 atgggagttg ccgccagatt ctgacatgga tctgaatctg attgagcagg cacccctgac 480 cgtggccgag aagctgcagc gcgacttcct ggtccactgg cgccgcgtga gtaaggcccc 540 ggaggccctc ttctttgttc agttcgagaa gggcgagtcc tacttccacc tccatattct 600 ggtggagacc acgggggtca aatccatggt gctgggccgc ttcctgagtc agattagcga 660 caagetggtg cagaceatet acegegggat egageegaee etgeecaaet ggttegeggt 720 gaccaagacg cgtaatggcg ccggaggggg gaacaaggtg gtggacgagt gctacatccc 780 caactacctc ctgcccaaga ctcagcccga gctgcagtgg gcgtggacta acatggagga 840 gtatataago gogtgtttaa acetggcoga gogcaaacgg ctcgtggcgc acgacetgac 900 ccacgtcagc cagacccagg agcagaacaa ggagaatctg aaccccaatt ctgacgcgcc 960 tgtcatccgg tcaaaaacct ccgcacgcta catggagctg gtcgggtggc tggtggaccg 1020 gggcatcaco teegagaage agtggateea ggaggaeeag geetegtaca teteetteaa 1080 cgccgcctcc aactcgcggt cccagatcaa ggccgctctg gacaatgccg gcaagatcat 1140 ggcgctgacc aaatccgcgc ccgactacct ggtaggcccc gctccgcccg ccgacattaa 1200 aaccaaccgc atttaccgca teetggaget gaacggetae gaecetgeet acgccggete 1260 cgtctttctc ggctgggccc agaaaaggtt cggaaaacgc aacaccatct ggctgtttgg 1320 gccggccacc acgggcaaga ccaacatcgc ggaagccatc gcccacgccg tgcccttcta 1380 cggctgcgtc aactggacca atgagaactt tcccttcaac gattgcgtcg acaagatggt 1440 gatctggtgg gaggagggca agatgacggc caaggtcgtg gagtccgcca aggccattct 1500

cggcggcagc aaggtgcgcg tggaccaaaa gtgcaagtcg tccgcccaga tcgatcccac 1560 ccccgtgatc gtcacctcca acaccaacat gtgcgccgtg attgacggga acagcaccac 1620 cttcgagcac cagcagccgt tgcaggaccg gatgttcaaa tttgaactca cccgccgtct 1680 ggagcatgac tttggcaagg tgacaaagca ggaagtcaaa gagttcttcc gctgggcgca 1740 ggatcacgtg accgaggtgg cgcatgagtt ctacgtcaga aagggtggag ccaacaacag 1800 accegecece gatgaegegg ataaaagega geecaagegg geetgeeeet eagtegegga 1860 tecategaeg teagaegegg aaggagetee ggtggaettt geegaeaggt accaaaacaa 1920 atgttetegt caegegggea tgetteagat getgttteee tgeaaaacat gegagagaat 1980 qaatcagaat ticaacatit gottoacgoa egggaccaga gactgitcag aaigtitcoc 2040 cggcgtgtca gaatctcaac cggtcgtcag aaagaggacg tatcggaaac tctgtgccat 2100 teateatetg etggggeggg etecegagat tgettgeteg geetgegate tggteaacgt 2160 qqatctqqat gactqtqttt ctqaqcaata aatqacttaa accaqqtatq gctqccqatq 2220 qttatcttcc agattggctc gaggacaacc tctctgaggg cattcggcag tggtgggact 2280 tgaaacctgg agccccgaaa cccaaagcca accagcaaaa gcaggacgac ggccggggtc 2340 tggtgcttcc tggctacaag tacctcggac ccttcaacgg actcgacaag ggggagcccg 2400 tcaacgegge ggatgeageg geeetegage acgacaagge etacgaceag cageteaaag 2460 egggtgacaa teegtacetg eggtataace acgeegacge egagtttcag gagegtetge 2520 aagaagatac gtcttttggg ggcaacctcg ggcgagcagt cttccaggcc aagaagaggg 2580 ttctcgaacc ttttggtctg gttgaggaag gtgctaagac ggctcctgga aagaaacgtc 2640 cggtagagca gtcgccacaa gagccagact cctcctcggg cattggcaag acaggccagc 2700 agecegetaa aaagagaete aattttggte agaetggega eteagagtea gteeeegaee 2760 cacaacctct cggagaacct ccagcaaccc ccgctgctgt gggacctact acaatggctt 2820 caggcggtgg cgcaccaatg gcagacaata acgaaggcgc cgacggagtg ggtaatgcct 2880 caggaaattg gcattgcgat tccacatggc tgggcgacag agtcatcacc accagcaccc 2940

4. (\*\*

gaacatgggc cttgcccacc tataacaacc acctctacaa gcaaatctcc agtgcttcaa 3000 cgggggccag caacgacaac cactacttcg gctacagcac cccctggggg tattttgatt 3060 tcaacagatt ccactgccat ttctcaccac gtgactggca gcgactcatc aacaacaatt 3120 ggggatteeg geccaagaga eteaaettea agetetteaa catecaagte aaggaggtea 3180 cgacgaatga tggcgtcacg accatcgcta ataaccttac cagcacggtt caagtcttgt 3240 eggactegga gtaccagtte eegtacgtee teggetetge geaceaggge tgeeteecte 3300 cgttcccggc ggacgtgttc atgattccgc agtacggcta cctaacgctc aacaatggca 3360 gecaggeagt gggaegetea teettttaet geetggaata ttteecateg eagatgetga 3420 gaacgggcaa taactttacc ttcagctaca ccttcgagga cgtgcctttc cacagcagct 3480 acgcgcacag ccagagcetg gaccggctga tgaatcetet catcgaccag tacetgtatt 3540 acctgaacag aactcacaat cagtccggaa gtgcccaaaa caaggacttg ctgtttagcc 3600 gtgggtetee agetggeatg tetgtteage ceaaaaactg getacetgga ecetgttace 3660 gqcagcagcg cgtttctaaa acaaaaacag acaacaacaa cagcaacttt acctqqactq 3720 gtgcttcaaa atataacctt aatggeegtg aatctataat caaccctggc actgctatgg 3780 cctcacacaa agacgacaaa gacaagttct ttcccatgag cggtgtcatg atttttggaa 3840 aggagagege eggagettea aacactgeat tggacaatgt catgateaca gaegaagagg 3900 aaatcaaagc cactaacccc gtggccaccg aaagatttgg gactgtggca gtcaatctcc 3960 agagcagcag cacagaccct gcgaccggag atgtgcatgt tatgggagcc ttacctggaa 4020 tggtgtggca agacagagac gtatacctgc agggtcctat ttgggccaaa attcctcaca 4080 eggatggaca ettteaceeg teteetetea tgggeggett tggaettaag caccegeete 4140 ctcagatcct catcaaaaac acgcctgttc ctgcgaatcc tccggcagag ttttcggcta 4200 caaagtttgc ttcattcatc acccagtatt ccacaggaca agtgagcgtg gagattgaat 4260 gggagctgca gaaagaaaac aycaaacgct ggaatcccga agtgcagtat acatctaact 4320 atgcaaaatc tgccaacgtt gatttcactg tggacaacaa tggactttat actgagcctc 4380

generating caccegitate eteacocyte contiguant gigitytaat caataaaccy 4440 gitaattegi gicagityaa etitigytete atgicettat tatettatet gigitaaccata 4500 geaacciggit acacattaac tigettagity egettegega ataccectag tigatygagit 4560 geecacteec tetatyeegi etegeteget egigityggee gigiagagia agetetyeegi 4620 tetgeggace titiggiteegi aggeecace gagegagega gegegeatag aggagtyge 4680 caa 4683

<210> 20

<211> 16

<212> DNA

<213> rep binding motif

<400> 20

gctcgctcgc tcgctg

## This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
D BLACK BORDERS
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
☐ SKEWED/SLANTED IMAGES
COLOR OR BLACK AND WHITE PHOTOGRAPHS
☐ GRAY SCALE DOCUMENTS
☐ LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
OTHER.

## IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.